

(19) World Intellectual Property Organization
International Bureau




(43) International Publication Date
15 March 2001 (15.03.2001)

PCT

(10) International Publication Number
WO 01/18225 A1

- (51) International Patent Classification⁷: C12N 15/90, A01K 67/027
- (21) International Application Number: PCT/US99/30078
- (22) International Filing Date:
16 December 1999 (16.12.1999)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/152,522 3 September 1999 (03.09.1999) US
- (71) Applicant: XENOGEN CORPORATION [US/US]; 860 Atlantic Avenue, Alameda, CA 94501 (US).
- (72) Inventor: ZHANG, Ning; Xenogen Corporation, 860 Atlantic Avenue, Alameda, CA 94501 (US).
- (74) Agent: SHOLTZ, Charles, K.; Xenogen Corporation, 860 Atlantic Avenue, Alameda, CA 94501 (US).
- (81) Designated States (*national*): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— With international search report.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: TARGETING CONSTRUCTS AND TRANSGENIC ANIMALS PRODUCED THEREWITH



(57) Abstract: The present invention teaches targeting constructs and methods of use thereof for creating transgenic animals in which at least one single-copy, non-essential gene is replaced with a reporter expression cassette, for example, a luciferase gene operably linked to a promoter heterologous to the single-copy, non-essential gene. Thus, the present invention provides novel methods and vector constructs useful for the generation of transgenic animals. The invention further includes methods of using these animals.

TARGETING CONSTRUCTS AND TRANSGENIC ANIMALS PRODUCED THEREWITH

5

TECHNICAL FIELD

This invention is in the field of molecular biology and medicine. More specifically, it relates to novel vector constructs and methods of use thereof for introducing heterologous polynucleotides into a host cell. Further, the invention relates to vector constructs and methods of use thereof to generate transgenic organisms, particularly transgenic mice.

BACKGROUND

In recent years, mouse geneticists have succeeded in creating transgenic animals by manipulating the genes of developing embryos and introducing foreign genes into these embryos. Once these genes have integrated into the genome of the recipient embryo, the resulting embryos or adult animals can be analyzed to determine the function of the gene.

Traditionally, transgenic mice have been generated through DNA microinjection approach. Such an approach leads to the creation of "founder" mice having at least one copy of the transgene randomly integrated into the genome. Because neither the copy number nor the integration sites can be controlled, transgenic mice generated by this method are genetically different from each other. The expression of transgenes in such transgenic mice are not uniform because of, for example, the difference in copy numbers of the transgene. Furthermore, chromosomal location of the transgene often affects the expression level of the transgene. For most of *in vivo* studies, particularly ones where it is desirable to compare levels of gene expression across different animals, it is important to have the mice with little or no genetic variation (i.e., isogenic mice) in order to reduce the systematic error.

One way of controlling copy number and the chromosomal integration site is by using targeting constructs to create transgenic animals. U.S. Patent Nos. 5,464,764 and 5,487,992 describe this type of transgenic animal in which a gene of interest is deleted or mutated sufficiently to disrupt its function. These "knock-out" animals are made by taking advantage of the phenomena of homologous recombination. (See, also U.S.

Patent Nos. 5,631,153 and 5,627,059). Using the "knock-out" approach, when the vector is introduced into the embryonic stem cell, a sequence becomes integrated into a target gene in the genome via homologous recombination. The integration disrupts the function of the target gene allows for examination of the phenotype resulting from the disruption of the gene. Neither of the above methods, however, provide a predictable approach for the generation of gene expression reporter transgenic animals which can be used for quantitative comparisons of gene expression across different animals transfected with the same construct, or for quantitative comparisons of the relative levels of expression of two or more different genes.

The present invention solves this and other problems by providing transgenic animals in which at least one single-copy, non-essential gene is replaced with a reporter expression cassette, for example, a luciferase gene operably linked to a heterologous promoter. Thus, the present invention provides novel methods and vector constructs useful for the generation of transgenic animals. The invention further includes methods of using these animals.

SUMMARY OF THE INVENTION

The transgenic animals described herein are useful, for example, when studying *in vivo* regulation of selected genes. Also described herein are methods of generating populations of substantially isogenic transgenic animals, as well as, vectors useful in these methods.

Accordingly, in one embodiment, the subject invention is directed to a transgenic, non-human mammal, for example, a rodent such as a mouse. The mammal comprises at least one single-copy, non-essential gene in its genome, wherein (i) at least a portion of at least one single-copy, non-essential gene is replaced by polynucleotide sequences heterologous to the gene, and (ii) the polynucleotide sequences comprise a first expression cassette which has been introduced into the mammal or an ancestor of the mammal, at an embryonic stage. The first expression cassette typically comprises a first selectable marker, a first transcriptional promoter element heterologous to the gene, and light-generating protein coding sequences. The light-generating protein coding sequences are operably linked to the promoter element.

The single-copy, non-essential gene may be selected, for example, from the group consisting of vitronectin, *fosB*, and galactin 3 and the first selectable marker may be selected from the group consisting of neomycin phosphotransferase II, xanthine-guanine phosphoribosyltransferase, hygromycin-B-phosphotransferase, chloramphenicol acetyltransferase, and adeninephosphoribosyl transferase.

In alternative embodiments, the first transcriptional promoter element is an inducible promoter, a repressible promoter, or a constitutive promoter, and may be selected from the group consisting of VEGF, VEGFR, and TIE2.

In additional embodiments, the transgenic, non-human mammal described above comprises a second single-copy, non-essential gene in its genome, wherein (i) at least a portion of the second single-copy, non-essential gene is replaced by polynucleotide sequences heterologous to the second gene, and (ii) the polynucleotide sequences comprise a second expression cassette which has been introduced into the mammal or an ancestor of the mammal, at an embryonic stage. The second expression cassette typically comprises a second selectable marker, a second transcriptional promoter element heterologous to the second gene, and light generating protein coding sequences. The light generating protein coding sequences are operably linked to the promoter element.

The first and second transcriptional promoter elements and selectable markers may be the same or different and the light generating protein in the first expression cassette can produce a different color of light relative to the light generating protein in the second expression cassette.

In yet a further embodiment, the invention is directed to a method of producing a transgenic, non-human mammal, such as a mouse. The mammal has at least one single-copy, non-essential gene in its genome. The method comprises

transfecting an embryonic stem cell of the mammal with a linear vector comprising

(a) a first selectable marker and a reporter expression cassette, the reporter expression cassette comprising a transcriptional promoter element operably linked to a light generating protein coding sequence, and

(b) targeting polynucleotide sequences homologous to a single-copy, non-essential gene in said mammal's genome, the targeting polynucleotide sequences

flanking (a), wherein (i) the length of the polynucleotide sequences are sufficient to facilitate homologous recombination between the vector and the single-copy, non-essential gene, and (ii) the transcriptional promoter element is heterologous to the single-copy, non-essential gene;

5 selecting embryonic stem cells which each have the first selectable marker and reporter expression cassette integrated into its genome;

 injecting the embryonic stem cells into a host embryo,

 implanting the embryo in a foster mother,

 maintaining the foster mother under conditions which allow production of an
10 offspring that is a transgenic, non-human mammal carrying the reporter expression cassette.

 In certain embodiments, the offspring is capable of germline transmission of the reporter expression cassette and the method may further comprise breeding the offspring with a mammal which is substantially isogenic with the embryonic stem cells, such that
15 the breeding yields transgenic F1 offspring carrying the reporter cassette. In particular embodiments, the method comprises breeding the first F1 offspring carrying the reporter cassette with a second F1 offspring carrying the reporter cassette, wherein the breeding yields transgenic F2 offspring carrying the reporter cassette.

 In additional embodiments, the embryonic stems cells may be derived from a
20 mouse having a dark coat color, the mammal substantially isogenic with the embryonic stem cells may have a light coat color, and/or the F2 offspring carrying the reporter cassette may have a light coat color. In particular embodiments, the embryonic stems cells are derived from a C57BL/6 mouse having a dark coat color, and the mammal substantially isogenic with the embryonic stem cells is a C57BL/6-Tyr C2j/+ mouse
25 having a light coat color.

 In still a further embodiment, the subject invention is directed to a vector for use in generating a transgenic non-human mammal, for example, a rodent such as a mouse. The mammal has at least one single-copy, non-essential gene in its genome. The vector comprises

30 (a) a first selectable marker and a reporter expression cassette, the reporter expression cassette comprising a transcriptional promoter element operably linked to a light generating protein coding sequence, and

(b) targeting polynucleotide sequences homologous to a single-copy, non-essential gene in the mammal's genome, the targeting polynucleotide sequences flanking (a), wherein (i) the length of the targeting polynucleotide sequences are sufficient to facilitate homologous recombination between the vector and the single-copy, non-essential gene, and (ii) the transcriptional promoter element is heterologous to the single-copy, non-essential gene.

In certain embodiments, the first selectable marker provides a positive selection and may be selected from the group consisting of neomycin phosphotransferase II, xanthine-guanine phosphoribosyltransferase, hygromycin-B-phosphotransferase, chloramphenicol acetyltransferase, and adeninephosphoribosyl transferase. Additionally, the transcriptional promoter element may be an inducible promoter, a repressible promoter, or a constitutive promoter, and may be selected from the group consisting of VEGF, VEGFR, and TIE2.

In alternative embodiments, the vector further comprises a second selectable marker and at least one target polynucleotide sequence is located between the second selectable marker and the first selectable marker. Additionally, the second selectable marker may provide a negative selection and may be selected from the group consisting of adenosine deaminase, thymidine kinase, and dihydrofolate reductase.

The vectors described above may be circular and may contain at least one restriction site whose cleavage results in a linear vector having the following arrangement of elements: target polynucleotide sequence - (a) - targeting polynucleotide sequences or target polynucleotide sequence - (a) - targeting polynucleotide sequences - (second selectable marker).

The coding sequences of the reporter expression cassette present in the vector may comprise codons that are optimal for expression in a host system into which the expression cassette is to be introduced. Additionally, the targeting polynucleotide sequences from single-copy, non-essential genes may be selected from the group consisting of vitronectin, *fosB*, and galactin 3.

The light-generating protein in the mammals and methods described above may be derived from either procaryotic or eucaryotic sources and, in particularly preferred embodiments, the light generating protein is a luciferase.

These and other embodiments of the present invention will be apparent to those of skill in the art in view of the teachings herein.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a schematic depicting construction of the pTK53 vector.

Polynucleotides encoding PGK-P, Neo and TK and 5' and 3' linkers are introduced into
5 a pKS backbone to produce the vector designated pTK53.

Figure 2 is schematic depicting construction of the pTK-LucR and pTK-LucYG vectors. For pTK-LucR, a polynucleotide encoding LucR is introduced into pTK53. Thus, the pTK-LucR construct contains the PGK-P gene, a neomycin (Neo^r) gene, a thymidine kinase (TK) gene and sequence encoding red luciferase (Luc-R). For pTK-
10 LucYG, a polynucleotide encoding LucYG is introduced into pTK53. Thus, the pTK-LucYG construct contains the PGK-P gene, a neomycin (Neo^r) gene, a thymidine kinase (TK) gene and a sequence encoding yellow-green luciferase (Luc-YG).

Figures 3A is a schematic depicting the vector pTKLR-Vn. Sequences homologous to the vitronectin gene are inserted into pTK-LucR such that they flank the
15 Neo^r gene and the Luc-R coding sequence. Figure 3B is a schematic depicting targeting of the linearized pTKLR-Vn vector to the vitronectin chromosomal locus. The VEGF promoter is cloned into the polylinkers between Neo and Luc-R. Upon homologous recombination, the Neo-VEGF-LucR transgene is inserted into the Vn gene. In the figure, (A) shows the targeting vector pTKLR-Vn and (B) shows the mouse vitronectin
20 gene. In the figure, Neo – neomycin resistance encoding sequences; TK – thymidine kinase encoding sequences; LucR – red luciferase from pGL3Red (Dr. Christopher Contag, Stanford University, Stanford, CA). Regions bearing Vn gene translational start and stop codons are indicated with arrows. Poly(A) sequences are placed upstream of the polylinker to prevent or minimize read-through translation. Figure 3C shows the
25 nucleotide sequence of vitronectin (SEQ ID NO:38).

Figure 4A is a schematic depicting the vector pTKLG-Fos. Sequences homologous to the FosB gene are inserted into pTK-LucYG such that they flank the Neo^r gene and the Luc-YG coding sequence. Figure 4B shows the nucleotide sequence of FosB (SEQ ID NO:39).

30 Figure 5A is a schematic depicting targeting of the linearized pTKLG-Fos vector to the FosB chromosomal locus. The VEGFR2 promoter is cloned into the polylinkers between Neo and Luc-YG. Upon homologous recombination, the Neo-VEGFR2-LucYG

transgene will be inserted into a sequence associated with production of FosB. In the figure, (A) shows the targeting vector, and (B) shows the mouse target gene. In the figure, Neo – neomycin resistance encoding sequences; TK – thymidine kinase encoding sequences; LucYG – yellow green luciferase from pGL3-control vector (Promega, Madison, WI). Regions bearing FosB gene translational start and stop codons are indicated with arrows. Poly(A) sequences are placed upstream of the polylinker to prevent or minimize read-through translation. Figure 5B is a schematic depicting targeting of the linearized pTKLG-Fos vector to the FosB chromosomal locus. The TIE2 promoter is cloned into the polylinkers between Neo and Luc-R. Upon homologous recombination, the Neo-Tie2-LucYG transgene is inserted into the FosB gene. In the figure, (A) shows the targeting vector, and (B) shows the mouse target gene. In the figure, Neo – neomycin resistance encoding sequences; TK – thymidine kinase encoding sequences; LucYG – yellow green luciferase from pGL3-control vector (Promega). Regions bearing FosB gene translational start and stop codons are indicated with arrows. Poly(A) sequences are placed upstream of the polylinker to prevent or minimize read-through translation.

Figure 6 depicts PCR conditions for genomic screening for promoters useful in exemplary targeting constructs of the present invention.

Figure 7 depicts generation of targeted transgenic mice using the targeting vectors described herein.

Figure 8 depicts of schematic representation of Southern blot analysis of homologous DNA recombination between pTKLG-Fos targeting vector and the FosB gene.

Figure 9 depicts generation of targeted transgenic mice, using the targeting vectors described herein, and crosses using such transgenics as well as their offspring (F1, first generation; F2, second generation).

Figure 10 depicts crosses using transgenic mice of the present invention to generate dual luciferase transgenic mice.

MODES FOR CARRYING OUT THE INVENTION

Throughout this application, various publications, patents, and published patent applications are referred to by an identifying citation. The disclosures of these publications, patents, and published patent specifications referenced in this application
5 are included to more fully describe the state of the art to which this invention pertains.

The practice of the present invention will employ, unless otherwise indicated, conventional techniques of molecular biology, microbiology, cell biology and recombinant DNA, which are within the skill of the art. *See, e.g.*, Sambrook, Fritsch, and Maniatis, MOLECULAR CLONING: A LABORATORY MANUAL, 2nd edition
10 (1989); CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, (F.M. Ausubel et al. eds., 1987); the series METHODS IN ENZYMOLOGY (Academic Press, Inc.); PCR 2: A PRACTICAL APPROACH (M.J. McPherson, B.D. Hames and G.R. Taylor eds., 1995) and ANIMAL CELL CULTURE (R.I. Freshney. Ed., 1987).

As used in this specification and the appended claims, the singular forms "a,"
15 "an" and "the" include plural references unless the content clearly dictates otherwise. Thus, for example, reference to "an antigen" includes a mixture of two or more such agents.

Definitions

20 As used herein, certain terms will have specific meanings.

The terms "nucleic acid molecule" and "polynucleotide" are used interchangeably to and refer to a polymeric form of nucleotides of any length, either deoxyribonucleotides or ribonucleotides, or analogs thereof. Polynucleotides may have any three-dimensional structure, and may perform any function, known or unknown.
25 Non-limiting examples of polynucleotides include a gene, a gene fragment, exons, introns, messenger RNA (mRNA), transfer RNA, ribosomal RNA, ribozymes, cDNA, recombinant polynucleotides, branched polynucleotides, plasmids, vectors, isolated DNA of any sequence, isolated RNA of any sequence, nucleic acid probes, and primers.

A polynucleotide is typically composed of a specific sequence of four nucleotide
30 bases: adenine (A); cytosine (C); guanine (G); and thymine (T) (uracil (U) for thymine (T) when the polynucleotide is RNA). Thus, the term polynucleotide sequence is the alphabetical representation of a polynucleotide molecule. This alphabetical

representation can be input into databases in a computer having a central processing unit and used for bioinformatics applications such as functional genomics and homology searching.

A "coding sequence" or a sequence which "encodes" a selected polypeptide, is a nucleic acid molecule which is transcribed (in the case of DNA) and translated (in the case of mRNA) into a polypeptide, for example, *in vivo* when placed under the control of appropriate regulatory sequences (or "control elements"). The boundaries of the coding sequence are typically determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxy) terminus. A coding sequence can include, but is not limited to, cDNA from viral, procaryotic or eucaryotic mRNA, genomic DNA sequences from viral or procaryotic DNA, and even synthetic DNA sequences. A transcription termination sequence may be located 3' to the coding sequence. Other "control elements" may also be associated with a coding sequence. A DNA sequence encoding a polypeptide can be optimized for expression in a selected cell by using the codons preferred by the selected cell to represent the DNA copy of the desired polypeptide coding sequence. "Encoded by" refers to a nucleic acid sequence which codes for a polypeptide sequence, wherein the polypeptide sequence or a portion thereof contains an amino acid sequence of at least 3 to 5 amino acids, more preferably at least 8 to 10 amino acids, and even more preferably at least 15 to 20 amino acids from a polypeptide encoded by the nucleic acid sequence. Also encompassed are polypeptide sequences which are immunologically identifiable with a polypeptide encoded by the sequence.

Typical "control elements", include, but are not limited to, transcription promoters, transcription enhancer elements, transcription termination signals, polyadenylation sequences (located 3' to the translation stop codon), sequences for optimization of initiation of translation (located 5' to the coding sequence), translation enhancing sequences, and translation termination sequences. Transcription promoters can include inducible promoters (where expression of a polynucleotide sequence operably linked to the promoter is induced by an analyte, cofactor, regulatory protein, etc.), repressible promoters (where expression of a polynucleotide sequence operably linked to the promoter is induced by an analyte, cofactor, regulatory protein, etc.), and constitutive promoters.

"Expression enhancing sequences" typically refer to control elements that improve transcription or translation of a polynucleotide relative to the expression level in the absence of such control elements (for example, promoters, promoter enhancers, enhancer elements, and translational enhancers (e.g., Shine and Delagarno sequences)).

5 "Purified polynucleotide" refers to a polynucleotide of interest or fragment thereof which is essentially free, e.g., contains less than about 50%, preferably less than about 70%, and more preferably less than about 90%, of the protein with which the polynucleotide is naturally associated. Techniques for purifying polynucleotides of interest are well-known in the art and include, for example, disruption of the cell
10 containing the polynucleotide with a chaotropic agent and separation of the polynucleotide(s) and proteins by ion-exchange chromatography, affinity chromatography and sedimentation according to density.

A "heterologous sequence" as used herein is typically refers to either (i) a nucleic acid sequence that is not normally found in the cell or organism of interest, or (ii) a
15 nucleic acid sequence introduced at a genomic site wherein the nucleic acid sequence does not normally occur in nature at that site. For example, a DNA sequence encoding a polypeptide can be obtained from yeast and introduced into a bacterial cell. In this case the yeast DNA sequence is "heterologous" to the native DNA of the bacterial cell. Alternatively, a promoter sequence from a Tie2 gene can be introduced into the genomic
20 location of a *fosB* gene. In this case the Tie2 promoter sequence is "heterologous" to the native *fosB* genomic sequence.

A "polypeptide" is used in it broadest sense to refer to a compound of two or more subunit amino acids, amino acid analogs, or other peptidomimetics. The subunits may be linked by peptide bonds or by other bonds, for example ester, ether, etc. As used
25 herein, the term "amino acid" refers to either natural and/or unnatural or synthetic amino acids, including glycine and both the D or L optical isomers, and amino acid analogs and peptidomimetics. A peptide of three or more amino acids is commonly called an oligopeptide if the peptide chain is short. If the peptide chain is long, the peptide is typically called a polypeptide or a protein.

30 "Operably linked" refers to an arrangement of elements wherein the components so described are configured so as to perform their usual function. Thus, a given promoter that is operably linked to a coding sequence (e.g., a reporter expression

cassette) is capable of effecting the expression of the coding sequence when the proper enzymes are present. The promoter or other control elements need not be contiguous with the coding sequence, so long as they function to direct the expression thereof. For example, intervening untranslated yet transcribed sequences can be present between the promoter sequence and the coding sequence and the promoter sequence can still be
5 considered "operably linked" to the coding sequence.

"Recombinant" as used herein to describe a nucleic acid molecule means a polynucleotide of genomic, cDNA, semisynthetic, or synthetic origin which, by virtue of its origin or manipulation: (1) is not associated with all or a portion of the polynucleotide
10 with which it is associated in nature; and/or (2) is linked to a polynucleotide other than that to which it is linked in nature. The term "recombinant" as used with respect to a protein or polypeptide means a polypeptide produced by expression of a recombinant polynucleotide. "Recombinant host cells," "host cells," "cells," "cell lines," "cell cultures," and other such terms denoting procaryotic microorganisms or eucaryotic cell
15 lines cultured as unicellular entities, are used interchangeably, and refer to cells which can be, or have been, used as recipients for recombinant vectors or other transfer DNA, and include the progeny of the original cell which has been transfected. It is understood that the progeny of a single parental cell may not necessarily be completely identical in morphology or in genomic or total DNA complement to the original parent, due to
20 accidental or deliberate mutation. Progeny of the parental cell which are sufficiently similar to the parent to be characterized by the relevant property, such as the presence of a nucleotide sequence encoding a desired peptide, are included in the progeny intended by this definition, and are covered by the above terms. An "isolated polynucleotide" molecule is a nucleic acid molecule separate and discrete from the whole organism with
25 which the molecule is found in nature; or a nucleic acid molecule devoid, in whole or part, of sequences normally associated with it in nature; or a sequence, as it exists in nature, but having heterologous sequences (as defined below) in association therewith.

Techniques for determining nucleic acid and amino acid "sequence identity" also are known in the art. Typically, such techniques include determining the nucleotide
30 sequence of the mRNA for a gene and/or determining the amino acid sequence encoded thereby, and comparing these sequences to a second nucleotide or amino acid sequence. In general, "identity" refers to an exact nucleotide-to-nucleotide or amino acid-to-amino

acid correspondence of two polynucleotides or polypeptide sequences, respectively. Two or more sequences (polynucleotide or amino acid) can be compared by determining their "percent identity." The percent identity of two sequences, whether nucleic acid or amino acid sequences, is the number of exact matches between two aligned sequences divided by the length of the shorter sequences and multiplied by 100. An approximate alignment for nucleic acid sequences is provided by the local homology algorithm of Smith and Waterman, Advances in Applied Mathematics 2:482-489 (1981). This algorithm can be applied to amino acid sequences by using the scoring matrix developed by Dayhoff, Atlas of Protein Sequences and Structure, M.O. Dayhoff ed., 5 suppl. 3:353-358, National Biomedical Research Foundation, Washington, D.C., USA, and normalized by Gribskov, Nucl. Acids Res. 14(6):6745-6763 (1986). An exemplary implementation of this algorithm to determine percent identity of a sequence is provided by the Genetics Computer Group (Madison, WI) in the "BestFit" utility application. The default parameters for this method are described in the Wisconsin Sequence Analysis Package Program Manual, Version 8 (1995) (available from Genetics Computer Group, Madison, WI). A preferred method of establishing percent identity in the context of the present invention is to use the MPSRCH package of programs copyrighted by the University of Edinburgh, developed by John F. Collins and Shane S. Sturrok, and distributed by IntelliGenetics, Inc. (Mountain View, CA). From this suite of packages the Smith-Waterman algorithm can be employed where default parameters are used for the scoring table (for example, gap open penalty of 12, gap extension penalty of one, and a gap of six). From the data generated the "Match" value reflects "sequence identity." Other suitable programs for calculating the percent identity or similarity between sequences are generally known in the art, for example, another alignment program is BLAST, used with default parameters. For example, BLASTN and BLASTP can be used using the following default parameters: genetic code = standard; filter = none; strand = both; cutoff = 60; expect = 10; Matrix = BLOSUM62; Descriptions = 50 sequences; sort by = HIGH SCORE; Databases = non-redundant, GenBank + EMBL + DDBJ + PDB + GenBank CDS translations + Swiss protein + Spupdate + PIR. Details of these programs can be found at the following internet address: <http://www.ncbi.nlm.gov/cgi-bin/BLAST>. When claiming sequences relative to sequences of the present invention, the desired degrees of sequence identity are at least

80%, 85-90%, preferably 92%, more preferably 95%, and even more preferably 98% sequence identity to the reference sequence (i.e., the sequences of the present invention).

Alternatively, the degree of sequence similarity between polynucleotides can be determined by hybridization of polynucleotides under conditions that form stable
5 duplexes between homologous regions, followed by digestion with single-stranded-specific nuclease(s), and size determination of the digested fragments. Two DNA, or two polypeptide sequences are "substantially homologous" to each other when the sequences exhibit at least about 80%-85%, preferably at least about 85%-90%, more preferably at least about 90%-95%, and most preferably at least about 95%-98%
10 sequence identity over a defined length of the molecules, as determined using the methods above. As used herein, substantially homologous also refers to sequences showing complete identity to the specified DNA or polypeptide sequence. DNA sequences that are substantially homologous can be identified in a Southern hybridization experiment under, for example, stringent conditions, as defined for that
15 particular system. Defining appropriate hybridization conditions is within the skill of the art. See, e.g., Sambrook et al., *supra*; *DNA Cloning, supra*; *Nucleic Acid Hybridization, supra*.

Two nucleic acid fragments are considered to "selectively hybridize" as described herein. The degree of sequence identity between two nucleic acid molecules affects the
20 efficiency and strength of hybridization events between such molecules. A partially identical nucleic acid sequence will at least partially inhibit a completely identical sequence from hybridizing to a target molecule. Inhibition of hybridization of the completely identical sequence can be assessed using hybridization assays that are well known in the art (e.g., Southern blot, Northern blot, solution hybridization, or the like,
25 see Sambrook, et al., *Molecular Cloning: A Laboratory Manual*, Second Edition, (1989) Cold Spring Harbor, N.Y.). Such assays can be conducted using varying degrees of selectivity, for example, using conditions varying from low to high stringency. If conditions of low stringency are employed, the absence of non-specific binding can be assessed using a secondary probe that lacks even a partial degree of sequence identity
30 (for example, a probe having less than about 30% sequence identity with the target molecule), such that, in the absence of non-specific binding events, the secondary probe will not hybridize to the target.

When utilizing a hybridization-based detection system, a nucleic acid probe is chosen that is complementary to a target nucleic acid sequence, and then by selection of appropriate conditions the probe and the target sequence "selectively hybridize," or bind, to each other to form a hybrid molecule. A nucleic acid molecule that is capable of

5 hybridizing selectively to a target sequence under "moderately stringent" typically hybridizes under conditions that allow detection of a target nucleic acid sequence of at least about 10-14 nucleotides in length having at least approximately 70% sequence identity with the sequence of the selected nucleic acid probe. Stringent hybridization conditions typically allow detection of target nucleic acid sequences of at least about 10-

10 14 nucleotides in length having a sequence identity of greater than about 90-95% with the sequence of the selected nucleic acid probe. Hybridization conditions useful for probe/target hybridization where the probe and target have a specific degree of sequence identity, can be determined as is known in the art (see, for example, Nucleic Acid Hybridization: A Practical Approach, editors B.D. Hames and S.J. Higgins, (1985)

15 Oxford; Washington, DC; IRL Press).

With respect to stringency conditions for hybridization, it is well known in the art that numerous equivalent conditions can be employed to establish a particular stringency by varying, for example, the following factors: the length and nature of probe and target sequences, base composition of the various sequences, concentrations of salts and other

20 hybridization solution components, the presence or absence of blocking agents in the hybridization solutions (e.g., formamide, dextran sulfate, and polyethylene glycol), hybridization reaction temperature and time parameters, as well as, varying wash conditions. The selection of a particular set of hybridization conditions is selected following standard methods in the art (see, for example, Sambrook, et al., Molecular

25 Cloning: A Laboratory Manual, Second Edition, (1989) Cold Spring Harbor, N.Y.).

A "vector" is capable of transferring gene sequences to target cells. Typically, "vector construct," "expression vector," and "gene transfer vector," mean any nucleic acid construct capable of directing the expression of a gene of interest and which can transfer gene sequences to target cells. Thus, the term includes cloning, and expression

30 vehicles, as well as integrating vectors.

"Nucleic acid expression vector" or "expression cassette" refers to an assembly which is capable of directing the expression of a sequence or gene of interest. The

nucleic acid expression vector includes a promoter which is operably linked to the sequences or gene(s) of interest. Other control elements may be present as well.

Expression cassettes described herein may be contained within a plasmid construct. In addition to the components of the expression cassette, the plasmid construct may also
5 include a bacterial origin of replication, one or more selectable markers, a signal which allows the plasmid construct to exist as single-stranded DNA (*e.g.*, a M13 origin of replication), a multiple cloning site, and a "mammalian" origin of replication (*e.g.*, a SV40 or adenovirus origin of replication).

An "expression cassette" comprises any nucleic acid construct capable of
10 directing the expression of a gene/coding sequence of interest. Such cassettes can be constructed into a "vector," "vector construct," "expression vector," or "gene transfer vector," in order to transfer the expression cassette into target cells. Thus, the term includes cloning and expression vehicles, as well as viral vectors.

"Luciferase," unless stated otherwise, includes prokaryotic and eukaryotic
15 luciferases, as well as variants possessing varied or altered optical properties, such as luciferases that produce different colors of light (*e.g.*, Kajiyama, N., and Nakano, E., *Protein Engineering* 4(6):691-693 (1991)).

"Light-generating" is defined as capable of generating light through a chemical reaction or through the absorption of radiation.

20 A "light generating protein" or "light-emitting protein" is a protein capable of generating light in the visible spectrum (between approximately 350 nm and 800 nm). Examples include bioluminescent proteins such as luciferases, *e.g.*, bacterial and firefly luciferases, as well as fluorescent proteins such as green fluorescent protein (GFP).

"Light" is defined herein, unless stated otherwise, as electromagnetic radiation
25 having a wavelength of between about 300 nm and about 1100 nm.

"Animal" as used herein typically refers to a non-human mammal, including, without limitation, farm animals such as cattle, sheep, pigs, goats and horses; domestic mammals such as dogs and cats; laboratory animals including rodents such as mice, rats and guinea pigs; birds, including domestic, wild and game birds such as chickens,
30 turkeys and other gallinaceous birds, ducks, geese, and the like. The term does not denote a particular age. Thus, both adult and newborn individuals are intended to be covered.

A "transgenic animal" refers to a genetically engineered animal or offspring of genetically engineered animals. A transgenic animal usually contains material from at least one unrelated organism, such as from a virus, plant, or other animal. The "non-human animals" of the invention include vertebrates such as rodents, non-human
5 primates, sheep, dogs, cows, amphibians, birds, fish, insects, reptiles, etc. The term "chimeric animal" is used to refer to animals in which the heterologous gene is found, or in which the heterologous gene is expressed in some but not all cells of the animal.

"Analyte" as used herein refers to any compound or substance whose effects (*e.g.*, induction or repression of a specific promoter) can be evaluated using the test
10 animals and methods of the present invention. Such analytes include, but are not limited to, chemical compounds, pharmaceutical compounds, polypeptides, peptides, polynucleotides, and polynucleotide analogs. Many organizations (*e.g.*, the National Institutes of Health, pharmaceutical and chemical corporations) have large libraries of chemical or biological compounds from natural or synthetic processes, or fermentation
15 broths or extracts. Such compounds/analytes can be employed in the practice of the present invention.

As used herein, the term "positive selection marker" refers to a gene encoding a product that enables only the cells that carry the gene to survive and/or grow under certain conditions. For example, plant and animal cells that express the introduced
20 neomycin resistance (Neo^r) gene are resistant to the compound G418. Cells that do not carry the Neo^r gene marker are killed by G418. Other positive selection markers will be known to those of skill in the art. Typically, positive selection markers encode products that can be readily assayed. Thus, positive selection markers can be used to determine whether a particular DNA construct has been introduced into a cell, organ or tissue.

25 "Negative selection marker" refers to gene encoding a product which can be used to selectively kill and/or inhibit growth of cells under certain conditions. Non-limiting examples of negative selection inserts include a herpes simplex virus (HSV)-thymidine kinase (TK) gene. Cells containing an active HSV-TK gene are incapable of growing in the presence of gangcylovir or similar agents. Thus, depending on the substrate, some
30 gene products can act as either positive or negative selection markers.

The term "homologous recombination" refers to the exchange of DNA fragments between two DNA molecules or chromatids at the site of essentially identical nucleotide

sequences. It is understood that substantially homologous sequences can accommodate insertions, deletions, and substitutions in the nucleotide sequence. Thus, linear sequences of nucleotides can be essentially identical even if some of the nucleotide residues do not precisely correspond or align (see, above).

5 A “knock-out” mutation refers to partial or complete loss of expression of at least a portion the target gene. Examples of knock-out mutations include, but are not limited to, gene-replacement by heterologous sequences, gene disruption by heterologous sequences, and deletion of essential elements of the gene (e.g., promoter region, portions of a coding sequence). A “knock-out” mutation is typically identified by the phenotype
10 generated by the mutation.

 A “single-copy gene” as used herein refers to a gene represented in an organism’s genome only by a single copy at a particular chromosomal locus. Accordingly, a diploid organism has two copies of the gene and both copies occur at the same chromosomal location.

15 A “gene” as used in the context of the present invention is a sequence of nucleotides in a genetic nucleic acid (chromosome, plasmid, etc.) with which a genetic function is associated. A gene is a hereditary unit, for example of an organism, comprising a polynucleotide sequence (e.g., a DNA sequence for mammals) that occupies a specific physical location (a “gene locus” or “genetic locus”) within the
20 genome of an organism. A gene can encode an expressed product, such as a polypeptide or a polynucleotide (e.g., tRNA). Alternatively, a gene may define a genomic location for a particular event/function, such as the binding of proteins and/or nucleic acids (e.g., phage attachment sites), wherein the gene does not encode an expressed product. Typically, a gene includes coding sequences, such as, polypeptide encoding sequences,
25 and non-coding sequences, such as, promoter sequences, poly-adenylation sequences, transcriptional regulatory sequences (e.g., enhancer sequences). Many eucaryotic genes have “exons” (coding sequences) interrupted by “introns” (non-coding sequences). In certain cases, a gene may share sequences with another gene(s) (e.g., overlapping genes).

 “Isogenic” means two or more organisms or cells that are considered to be
30 genetically identical. “Substantially isogenic” means two or more organisms or cells wherein, at the majority of genetic loci (e.g., greater than 99.000%, preferably more than 99.900%, more preferably greater than 99.990%, even more preferably greater than

99.999%), there exists genetic identity between the organisms or cells being compared. In the context of the present invention, two organisms (for example, mice) are considered to be "substantially isogenic" if, for example, inserted transgenes are the primary differences between the genetic make-up of the mice being compared. Further, if, for example, the genetic backgrounds of the mice being compared are the same with the exception that one of the mice has one or several defined mutation(s) (for example, affecting coat color), then these mice are considered to be substantially isogenic. An example of two strains of substantially isogenic mice are C57BL/6 and C57BL/6-Tyr C2j/+.

10 A "pseudogene" as used herein, refers to a type of gene sequence found in the genomes, typically, of eucaryotes, where the sequence closely resembles a known functional gene, but differs in that the pseudogene is non-functional. For example, the pseudogene sequence may contain several stop codons in what would correspond to an open reading frame in the functional gene. Pseudogenes can also have deletions or
15 insertions relative to their corresponding functional gene. If, for example, in a genome there is a functional gene and a related pseudogene, the functional gene is considered to be a single-copy gene (accordingly, the pseudogene is considered to be single-copy as well).

A "non-essential gene" refers to a gene whose deletion, disruption, elimination, reduction of gene function, or mutation is non-lethal, and does not obviously adversely affect the organisms' ability to mature and reproduce. A "non-essential gene with no phenotype" refers to a non-essential gene whose deletion, disruption, elimination, reduction of gene function or mutation has no deleterious effect on the organism. Typically there are no phenotypically reflected gene dosage effects associated with
25 modification of a non-essential gene with no phenotype -- for example, deletion, disruption or mutation of both copies of a non-essential gene with no phenotype in a diploid organism has essentially the same effect as deletion, disruption, or mutation of one of the two copies present in the diploid organism. In the context of the present invention, a non-essential gene is typically one whose function has been eliminated (e.g.,
30 by a deletion mutation) and such elimination of function was non-lethal and the organism developed, matured, and was able to reproduce.

The "native sequence" or "wild-type sequence" of a gene is the polynucleotide sequence that comprises the genetic locus corresponding to the gene, e.g., all regulatory and open-reading frame coding sequences required for expression of a completely functional gene product as they are present in the wild-type genome of an organism. The native sequence of a gene can include, for example, transcriptional promoter sequences, translation enhancing sequences, introns, exons, and poly-A processing signal sites. It is noted that in the general population, wild-type genes may include multiple prevalent versions that contain alterations in sequence relative to each other and yet do not cause a discernible pathological effect. These variations are designated "polymorphisms" or "allelic variations."

By "replacement sequence" is meant a polynucleotide sequence that is substituted for at least a portion of the native or wild-type sequence of a gene.

"Linear vector" or "linearized vector," as used herein, is a vector having two ends. For example, circular vectors, such as plasmids, can be linearized by digestion with a restriction endonuclease that cuts at a single site in the plasmid. Preferably, the targeting vectors described herein are linearized such that the ends are not within the targeting sequences.

General Overview

The present invention relates to vector constructs and methods of creating transgenic animals to be used, for example, as test systems. Methods of using the animals of the present invention include, but are not limited to, using these animals for studies involving tumor growth and other disease conditions. In the practice of the present invention, transgenic, non-human mammals are constructed where a single-copy, non-essential gene is replaced by a reporter expression cassette, preferably a gene encoding a light-generating protein, such as a luciferase-encoding gene, operably linked to a promoter. A variety of promoters are useful in the practice of the present invention, for example, promoters derived from genes associated with tumorigenesis or angiogenesis. Thus, an exemplary promoter can be one that is associated with proteins induced during tumorigenesis, for instance in the presence of tumor generating compounds or of tumors themselves. In this way, expression of the reporter cassette is induced in the animal when, for example, tumors are present, and progression of the

tumor can be evaluated by non-invasive imaging methods using the whole animal. Another exemplary promoter is one that is derived from a gene associated with angiogenesis. Because the promoter is linked to a reporter such as luciferase, non-invasive monitoring of the progression of angiogenesis is possible.

5 Various forms of the different embodiments of the invention, described herein, may be combined.

Non-invasive imaging and/or detecting of light-emitting conjugates in mammalian subjects was described in U.S. Patent No. 5,650,135, by Contag, et al., issued 22 July 1997. This imaging technology can be used in the practice of the present invention in view of the teachings of the present specification. In the imaging method, the conjugates contain a biocompatible entity and a light-generating moiety. Biocompatible entities include, but are not limited to, small molecules such as cyclic organic molecules; macromolecules such as proteins; microorganisms such as viruses, bacteria, yeast and fungi; eukaryotic cells; all types of pathogens and pathogenic substances; and particles such as beads and liposomes. In another aspect, biocompatible entities may be all or some of the cells that constitute the mammalian subject being imaged, for example, cells carrying the vector constructs of the present invention expressing a reporter expression cassette.

Light-emitting capability is conferred on the biocompatible entities by the conjugation of a light-generating moiety. Such moieties include fluorescent molecules, fluorescent proteins, enzymatic reactions giving off photons and luminescent substances, such as bioluminescent proteins. In the context of the present invention, light emitting capability is typically conferred on target cells by having at least one copy of a light-generating protein, *e.g.*, a luciferase, present. In preferred embodiments, luciferase is operably linked to appropriate control elements which can facilitate expression of a polypeptide having luciferase activity. Substrates of luciferase can be endogenous to the cell or applied to the cell or system (*e.g.*, injection into a transgenic mouse, having cells carrying a luciferase construct, of a suitable substrate for the luciferase, for example, luciferin). The conjugation may involve a chemical coupling step, genetic engineering of a fusion protein, or the transformation of a cell, microorganism or animal to express a light-generating protein.

Targeting Constructs

The targeting cassettes described herein typically include the following components: (1) a suitable vector backbone; (2) a polynucleotide encoding a light generating protein (3) a promoter operably linked to the light generating protein-
5 encoding gene, wherein the promoter is heterologous to the light generating protein coding sequences; (4) a sequence encoding a positive selection marker; (5) insertion sites flanking the sequence encoding the positive selection marker and the polynucleotide encoding a light generating protein gene, for insertion of sequences which target a single-copy, non-essential chromosomal gene; and, optionally, (6) a sequence encoding a
10 negative selection marker. Exemplary targeting constructs are shown in Figures 3B, 5A and 5B and described in Examples 1-3.

Suitable vector backbones generally include an F1 origin of replication; a colE1 plasmid-derived origin of replication; polyadenylation sequence(s); sequences encoding antibiotic resistance (*e.g.*, ampicillin resistance) and other regulatory or control elements.
15 Non-limiting examples of appropriate backbones include: pBluescriptSK (Stratagene, La Jolla, CA); pBluescriptKS (Stratagene, La Jolla, CA) and other commercially available vectors.

In one aspect of the invention the light generating protein is luciferase. Luciferase coding sequences useful in the practice of the present invention include, but
20 are not limited to, sequences obtained from *lux* genes (procaryotic genes encoding a luciferase activity) and *luc* genes (eucaryotic genes encoding a luciferase activity). A variety of luciferase encoding genes have been identified including, but not limited to, the following: B.A. Sherf and K.V. Wood, U.S. Patent No. 5,670,356, issued 23 September 1997; Kazami, J., et al., U.S. Patent No. 5,604,123, issued 18 February 1997;
25 S. Zenno, et al, U.S. Patent No. 5,618,722; K.V. Wood, U.S. Patent No. 5,650,289, issued 22 July 1997; K.V. Wood, U.S. Patent No. 5,641,641, issued 24 June 1997; N. Kajiyama and E. Nakano, U.S. Patent No. 5,229,285, issued 20 July 1993; M.J. Cormier and W.W. Lorenz, U.S. Patent No. 5,292,658, issued 8 March 1994; M.J. Cormier and W.W. Lorenz, U.S. Patent No. 5,418,155, issued 23 May 1995; de Wet, J.R., et al,
30 *Molec. Cell. Biol.* 7:725-737, 1987; Tatsumi, H.N., et al, *Biochim. Biophys. Acta* 1131:161-165, 1992; and Wood, K.V., et al, *Science* 244:700-702, 1989. Eukaryotic luciferase catalyzes a reaction using luciferin as a luminescent substrate to produce light,

whereas prokaryotic luciferase catalyzes a reaction using an aldehyde as a luminescent substrate to produce light.

Wild-type firefly luciferases typically have an emission maxima at about 550 nm. Numerous variants with differing emission maxima have also been studied. For example, Kajiyama and Nakano (*Protein Eng.* 4(6):691-693, 1991; U.S. Patent No. 5,330,906, issued 19 July 1994) teach five variant firefly luciferases generated by single amino acid changes to the *Luciola cruciata* luciferase coding sequence. The variants have emission peaks of 558 nm, 595 nm, 607 nm, 609 nm and 612 nm. A yellow-green luciferase with an emission peak of about 540 nm is commercially available from Promega, Madison, WI under the name pGL3. A red luciferase with an emission peak of about 610 nm is described, for example, in Contag et al. (1998) *Nat. Med.* 4:245-247 and Kajiyama et al. (1991) *Prot. Eng.* 4:691-693.

Positive selection markers include any gene which a product that can be readily assayed. Examples include, but are not limited to, a hprt gene (Littlefield, J. W., Science 145:709-710 (1964)), a xanthine-guanine phosphoribosyltransferase (gpt) gene, or an adenosine phosphoribosyltransferase (aprt) gene (Sambrook et al., *supra*), a thymidine kinase gene (i.e. "TK") and especially the TK gene of herpes simplex virus (Giphart-Gassler, M. et al., *Mutat. Res.* 214:223-232 (1989)), a nptII gene (Thomas, K. R. et al., *Cell* 51:503-512 (1987); Mansour, S. L. et al., *Nature* 336:348-352 (1988)), or other genes which confer resistance to amino acid or nucleoside analogues, or antibiotics, etc, for example, gene sequences which encode enzymes such as dihydrofolate reductase (DHFR) enzyme, adenosine deaminase (ADA), asparagine synthetase (AS), hygromycin B phosphotransferase, or a CAD enzyme (carbamyl phosphate synthetase, aspartate transcarbamylase, and dihydroorotase). Addition of the appropriate substrate of the positive selection marker can be used to determine if the product of the positive selection marker is expressed, for example cells which do not express the positive selection marker nptII, are killed when exposed to the substrate G418 (Gibco BRL Life Technology, Gaithersburg, MD).

The targeting vector typically contains insertion sites for inserting targeting sequences (e.g., sequences that are substantially homologous to the target sequences in the host genome where integration of the targeting vector/expression cassette is desired). These insertion sites are preferably included such that there are two sites, one site on

either side of the sequences encoding the positive selection marker, light generating protein (e.g., luciferase) and the promoter. Insertion sites are, for example, restriction endonuclease recognition sites, and can, for example, represent unique restriction sites. In this way, the vector can be digested with the appropriate enzymes and the targeting
5 sequences ligated into the vector.

Optionally, the targeting construct can contain a polynucleotide encoding a negative selection marker. Suitable negative selection markers include, but are not limited to, HSV-tk (see, e.g., Majzoub et al. (1996) *New Engl. J. Med.* 334:904-907 and U.S. Patent No. 5,464,764), as well as genes encoding various toxins including the
10 diphtheria toxin, the tetanus toxin, the cholera toxin and the pertussis toxin. A further negative selection marker gene is the hypoxanthine-guanine phosphoribosyl transferase (HPRT) gene for negative selection in 6-thioguanine.

Exemplary promoters and single-copy, non-essential genes for use in the vector constructs and methods of the present invention are described below.

15

Promoters

The targeting constructs and transgenic animals described herein contain a sequence encoding a light generating protein, e.g., luciferase, gene operably linked to a promoter. The promoter may be from the same species as the transgenic animal (e.g.,
20 mouse promoter used in construct to make transgenic mouse) or from a different species (e.g., human promoter used in construct to make transgenic mouse). The promoter can be derived from any gene of interest. In one embodiment of the present invention, the promoter is derived from a gene whose expression is induced during angiogenesis, for example pathogenic angiogenesis like tumor development. Thus, when a tumor begins
25 to develop in a transgenic animal carrying a vector construct of the present invention, the promoter is induced and the animal expresses luciferase, which can then be monitored *in vivo*.

Exemplary promoters for use in the present invention are selected such that they are functional in a cell type and/or animal into which they are being introduced.

30 Exemplary promoters include, but are not limited to, promoters obtained from the following mouse genes: vascular endothelial growth factor (VEGF) (VEGF promoter described in U.S. Patent No. 5,916,763; Shima et al. (1996) *J. Bio. Chem.* 271:3877-

3883; sequence available on NCBI under accession number U41383); VEGFR2, also known as Flk-1, (VEGFR-2 promoter described, for example, in Röncke et al. (1996) *Circ. Res.* 79:277-285; Patterson et al. (1995) *J. Bio. Chem.* 270:23111-23118; Kappel et al. (1999) *Blood* 93:4282-4292; sequence available as accession number X89777 of NCBI database); Tie2, also known as Tek (Tie2 promoter described, for example, in Fadel et al. (1998) *Biochem. J.* 338:335-343; Schlaeger et al. (1995) *Develop.* 121:1089-1098; Schlager et al. (1997) *PNAS USA* 94:3058-3063). VEGF is a specific mitogen for EC *in vitro* and a potent angiogenic factor *in vivo*. In a tumorigenesis study, it was shown that VEGF was critical for the initial subcutaneous growth of T-47D breast carcinoma cells transplanted into nude mice, whereas other angiogenic factors, such as, bFGF can compensate for the loss of VEGF after the tumors have reached a certain size (Yoshiji, H., et al., 1997 *Cancer Research* 57: 3924-28). VEGF is a major mediator of aberrant EC proliferation and vascular permeability in a variety of human pathologic situation, such as, tumor angiogenesis, diabetic retinopathy and rheumatoid arthritis (Benjamin LE, et al., 1997 *PNAS* 94: 8761-66; Soker, S., et al., 1998 *Cell* 92: 735-745). VEGF is synthesized by tumor cells *in vivo* and accumulates in nearby blood vessels. Because leaky tumor vessels initiate a cascade of events, which include plasma extravasation and which lead ultimately to angiogenesis and tumor stroma formation, VEGF plays a pivotal role in promoting tumor growth (Dvorak, H.F., et al., 1991 *J Exp Med* 174:1275-8). VEGF expression was upregulated by hypoxia (Shweiki, D., et al., 1992 *Nature* 359: 843-5). VEGF is also upregulated by overexpression of v-Src oncogene (Mukhopadhyay, D., et al., 1995 *Cancer Res.* 15: 6161-5), c-SRC (Mukhopadhyay, D., et al., 1995 *Nature* 375: 577-81), and mutant ras oncogene (Plate, K.H., et al., 1992 *Nature* 359: 845-8). The tumor suppressor p53 downregulates VEGF expression (Mukhopadhyay, D., et al., 1995 *Cancer Res.* 15: 6161-5).

A number of cytokines and growth factors, including PGF and TPA (Grugel, S., et al., 1995 *J. Biological Chem.* 270: 25915-9), EGF, TGF- β , IL-1, IL-6 induce VEGF mRNA expression in certain type of cells (Ferrara, N., et al., 1997 *Endocr. Rev.* 18: 4-25). Kaposi's sarcoma-associated herpesvirus (KSHV) encoded a G-protein-coupled receptor, a homolog of IL-8 receptor, can activate JNK/SAPK and p38MAPK and increase VEGF production, thus causing cell transformation and tumorigenicity (Bais, C., et al., *Nature* 1998 391:86-9). VEGF overexpression in skin of transgenic mice

induces angiogenesis, vascular hyperpermeability and accelerated tumor development (Larcher, F., et al., *Oncogene* 1998 17:303-11).

VEGF-B (cDNA sequences available on databases) is a mitogen for EC and may be involved in angiogenesis in muscle and heart (Olofsson, B., et al., 1996 *Proc Natl Acad Sci U S A* 93:2576-81). Shown *in vitro*, binding of VEGF-B to its receptor VEGFR-1 leads to increased expression and activity of urokinase type plasminogen activator and plasminogen activator inhibitor, suggesting a role for VEGF-B in the regulation of extracellular matrix degradation, cell adhesion, and migration (Olofsson, B., et al., 1998 *Proc Natl Acad Sci U S A* 95:11709-14).

VEGF-C (see, e.g., U.S. Patent No. 5,916,763 and Shima et al., *supra*) may regulate angiogenesis of lymphatic vasculature, as suggested by the pattern of VEGF-C expression in mouse embryos (Kukk, E., et al., 1996 *Development* 122: 3829-37). Although VEGF-C is also a ligand for VEGFR-2, the functional significance of this potential interaction is unknown. Overexpression of VEGF-C in the skin of transgenic mice resulted in lymphatic, but not vascular, endothelial proliferation and vessel enlargement, suggesting the major function of VEGF-C is through VEGFR-3 rather than VEGFR-2 (Jeltsch M, et al., 1997 *Science* 276:1423-5). Shown by the CAM assay, VEGF and VEGF-C are specific angiogenic and lymphangiogenic growth factors, respectively (Oh, S.J., et al., (1997) *Devel. Biol.* 188: 96-109). VEGF-C overexpression in the skin of transgenic mice resulted in lymphatic, but not vascular, endothelial proliferation and vessel enlargement (Jeltsch M, et al., 1997 *Science* 276:1423-5).

VEGF-D (cDNA sequences available on databases) is a mitogen for EC. Given that VEGF-D can also activate VEGFR-3, it is possible that VEGF-D could be involved in the regulation of growth and/or differentiation of lymphatic endothelium (Achen, M.G., et al., 1998 *Proc Natl Acad Sci U S A* 95: 548-53). VEGF-D is induced by transcription factor c-Fos in mouse (Orlandini, M., 1996 *PNAS* 93: 11675-80).

VEGFR-1 signaling pathway may regulate normal endothelial cell-cell or cell matrix interactions during vascular development, as suggested by the knockout study (Fong, G.H., et al., 1995 *Nature* 376: 65-69). Although VEGFR-1 has a higher affinity to VEGF than VEGFR-2, it does not transduce the mitogenic signals of VEGF in ECs (Soker, S., et al., 1998 *Cell* 92: 735-745). VEGFR-2 (see, e.g., Röncke et al., Patterson et al., Kappel et al. (1999), *supra*) appears to be the major transducer of VEGF signals in

EC that result in chemotaxis, mitogenicity and gross morphological changes in target cells (Soker, S., et al., 1998 *Cell* 92: 735-745). VEGFR-3 has an essential role in the development of the embryonic cardiovascular system before the emergence of lymphatic vessels, as shown by the knockout study (Dumont, D.J., et al., 1998 *Science* 282: 946-949). Neuropillin-1 (see, e.g., Soker et al. (1998) *Cell* 92:735-745) is a receptor for VEGF165. It can enhance the binding of VEGF165 to VEGFR-2 and VEGF165 mediated chemotaxis (Soker, S., et al., 1998 *Cell* 92: 735-745). Neuropillin1 overexpression in transgenic mice resulted in embryonic lethality. The embryos possessed excess capillaries and blood vessels. Dilated vessels and hemorrhage were also observed (Kitsukawa, T., et al., 1995 *Development* 121: 4309-18).

Further promoters of interest include, but are not limited to, the following. Ang2 is expressed only at predominant vascular remodeling sites, such as ovary, placenta, uterus (Maisonpierre, P.C., et al., 1997 *Science* 277: 55-60). In glioblastoma angiogenesis, Ang2 is found to be expressed in endothelial cells of small blood vessel and capillaries while Ang1 is expressed in glioblastoma tumor cells (Stratmann, A., 1998 *Am J Pathol* 153: 1459-66). Ang2 is up-regulated in bovine microvascular endothelial by VEGF, bFGF, cytokines, hypoxia (Mandriota, S.J., 1998 *Circ Res* 83: 852-9). Ang2 transgenic overexpression disrupts angiogenesis, and is embryonic lethal (Maisonpierre, P.C., et al., 1997 *Science* 277: 55-60). Ang1 is widely expressed, less abundant in heart and liver (Maisonpierre, P.C., et al., 1997 *Science* 277: 55-60). Ang1 is expressed in mesenchymal cells and may up-regulate the expression of Tie2 in the endothelial cells (Suri, C., et al., 1996 *Cell* 87: 1171-1180). Ang1 overexpression in the skin of transgenic mice produces larger, more numerous, and more highly branched vessels (Suri, C., et al., *Science* 1998 282:468-71). Tie2 (see, e.g., Fadel et al.; Schlaeger et al. (1995), and Schlaeger et al. (1997), *supra*) is endothelial cell specific, up-regulated during wound healing, follicle maturation (Puri, M.C., et al., 1995 *EMBO J* 14: 5884-91) and pathologic angiogenesis (Kaipainen, A., 1994 *Cancer Research* 54: 6571-77), such as, glioblastoma (Stratmann, A., 1998 *Am J Pathol* 153: 1459-66). Tie2 is also expressed in non-proliferating adult endothelium and endothelial cell lines (Dumont, D.J., et al. (1994) *Genes & Develop.* 8:1897-1909). A Tie2 activating mutation causes vascular dysmorphogenesis (Vikkula M, et al., 1996 *Cell* 87: 1181-1190). Tie2 mutant overexpression in transgenic mice is embryonic lethal (Dumont, D.J., et al., *supra*).

Other promoters useful in the practice of the present invention include, by way of example, promoters derived from the sequences encoding the following polypeptide products: PTEN (dual specificity phosphatase); BAI (brain-specific angiogenesis inhibitor); KAI1 (KANGAI 1); catenin beta-1 (cadherin-associated protein, beta); COX2 (PTGS2 cyclooxygenase 2, a.k.a. prostaglandin-endoperoxide synthase 2); MMP2 (72 kDa Type IV-A collagenase); MMP9 (92 kDa type IV-B collagenase); TIMP2 (tissue inhibitor of metalloproteinase 2); and TIMP3 (tissue inhibitor of metalloproteinase 3).

PTEN is a tumor suppressor gene and encodes a protein of 403 amino acids. (Li et al. (1997) *Science* 275:1943-1946; DiCristofano et al. (1998) *Nature Genet.* 19:348-355). Overexpression of PTEN has been shown to inhibit cell migration and it is postulated that this protein may function as a tumor suppressor by negatively regulating cell interactions with the extracellular matrix or by negatively regulating the PI3K/PKB/Akt signaling pathway. (Tamura et al. (1998) *Science* 280:1614-1617; Stambolic et al. (1998) *Cell* 95:29-29). Mutations in PTEN have been detected in cancer cell lines and in the germline of patients having Cowden disease, Lhermitte-Duclos disease and Bannayan-Zonana syndrome (diseases and syndromes which are characterized by hyperplastic/dysplastic changes in the prostate, skin and colon and which are associated with an increased risk of certain cancers, for example, breast cancer, prostate cancer and colon cancer). (Marsh et al. (1998) *Hum. Molec. Genet.* 7:507-515; Marsh et al. (1998) *J. Med. Genet.* 35:881-885; Nelen et al. (1997) *Hum. Molec. Genet.* 6:1383-1387).

BAI1 protein is predicted to be 1,584 amino acids in length and includes an extracellular domain, an intracellular domain and a 7-span transmembrane region similar to that of the secretin receptor. (Nishimori et al. (1997) *Oncogene* 15:245-2150). The extracellular region of BAI1 has a single Arg-Gly-Asp (RGD) motif recognized by integrins and also has five sequences corresponding to the thrombospondin type I (accession number 188060) repeats that can inhibit angiogenesis includes by basic fibroblast growth factor (bFGF, accession number 134920). Shiratsuchi et al. (1997) *Cytogenet. Cell Genet.* 79:103-108, cloned 2 other brain-specific angiogenesis inhibiting genes, designated BAI2 (accession number 602683) and BAI 3 (accession number 602684). Thus, it is postulated that members of this gene family may play a role in suppression of glioblastoma.

KAI1 encodes a 267 amino acid protein which is a member of the leukocyte surface glycoprotein family. The protein has 4 hydrophobic transmembrane domains and 1 large extracellular hydrophilic domain with three potential N-glycosylation sites. (Dong et al. (1995) *Science* 268:884-886). Molecular analysis of KAI1 is described, for example, in Dong et al. (1997) *Genomics* 41:25-32. KAI1 is a tumor metastasis suppressor gene that is capable of inhibiting the metastatic process in experimental animals. Expression of KAI1 is downregulated during tumor progression of prostate, breast, lung, bladder and pancreatic cancers in humans, apparently at the transcriptional or posttranscriptional level. Mashimo et al. (1998) *PNAS USA* 95:11307-11311, found that the tumor suppressor gene p53 can directly inactivate the KAI1 gene by interacting with the region 5' to the coding sequence, suggesting a direct relationship between p53 and KAI1.

Catenin beta-1 is an adherens junction (AJ) protein, which are critical for establishing and maintaining epithelial cell layers, for instance during embryogenesis, wound healing and tumor cell metastasis. Molecular analysis, including description of sequence homology to plakoglobin (accession number 173325), homology to the drosophila gene "armadillo" and interactions with Lef1/Tcf DNA binding proteins, is described, for example, in Nollet et al. (1996) *Genomics* 32:413-424; McCrea et al. (1991) *Science* 254:1359-1361 and Korinek et al. (1997) *Science* 275:1784-1787. In addition, studies by Korinek et al., *supra* and Morin et al. (1997) *Science* 275:1787-1790, have indicated that APC (accession number 175100) negatively regulates catenin beta and that regulation of this protein is critical to the tumor suppressive effect of APC. Abnormally high levels of beta-catenin have been detected in certain human melanoma cell lines. (Rubinfeld et al. (1997) *Science* 275:1790-1792. Koch et al. (1999) *Cancer Res.* 59:269-273 report that childhood hepatoblastomas frequently carry a mutated degradation targeting box of the beta-catenin gene. Transgenic mice which express catenin beta under the control of an epidermal promoter undergo de novo hair morphogenesis and eventually these animals develop two types of tumors -- epithelioid cysts and trichofolliculomas. Gat et al. (1998) *Cell* 95:605-614.

COX2 encodes a cyclooxygenase and is a key regulator of prostaglandin synthesis. (Hla et al. (1992) *PNAS USA* 89:7384-7388; Jones et al. (1993) *J. Biol. Chem.* 268:9049-9054). In particular, COX2 is generally considered to be a mediator of

inflammation and overexpression of COX2 in rat epithelial cells results in elevated levels of E-cadherin and Bcl2. (Tsuji & DuBois (1995) *Cell* 83:493-501). In co-cultures of endothelial cells and colon carcinoma cells, cells that overexpress COX2 produce prostaglandins, proangiogenic factors and stimulate both endothelial migration and tube formation. (Tsuji et al. (1998) *Cell* 93:705-716). Experiments conducted using APC knock-out mice have demonstrated that animals homozygous for a disrupted COX2 locus develop significantly more adenomatous polyps. (Oshima et al. (1996) *Cell* 87:803-809). COX-2 "knock out" mice develop severe nephropathy, are susceptible to peritonitis, exhibit reduced arachidonic acid-induced inflammation and exhibit reduced indomethacin-induced gastric ulceration. (Morham et al. (1995) *Cell* 83:473-482; Langenbach et al. (1995) *Cell* 83:483-492). Female mice that are deficient in cyclooxygenase 2 exhibit multiple reproductive failures. (Lim et al. (1997) *Cell* 91:197-208).

MMP2 is a metalloproteinase that specifically cleaves type IV collagen. A C-terminal fragment of MMP2, termed PEX, prevents normal binding to α -V/ β -3 and disrupts angiogenesis and tumor growth. (Brooks et al. (1998) *Cell* 92:391-400).

MMP9 is a collagenase secreted from normal skin fibroblasts. MMP9 null mice exhibit an abnormal pattern of skeletal growth plate vascularization and ossification. (Vu et al. (1998) *Cell* 93:411-422).

TIMP2 is a collagenase and appears to play a major role in modulating the activity of interstitial collagenase and a number of connective tissue metalloendoproteases. (Stetler-Stevenson et al. (1989) *J. Biol. Chem.* 264:17372-17378). Unlike TIMP1 and TIMP3, TIMP2 is not upregulated by TPA or TGF- β . (Hammani et al. (1996) *J. Biol. Chem.* 271:25498-25505).

TIMP3 (Wilde et al. (1994) *DNA Cell Biol.* 13:711-718) is localized in the extracellular matrix in both its glycosylated and unglycosylated forms. Studies of mutant TIMP3 proteins have demonstrated that C-terminal truncations do not bind to the extracellular matrix. (Langton et al. (1998) *J. Biol. Chem.* 273:16778-16781).

As one of skill in the art will appreciate in view of the teachings of the present specification, promoter sequences can be easily derived and isolated from known polypeptide sequences or from cDNA or genomic sequences, using method known in the art in view of the teachings herein. An exemplary method of isolating promoter

sequences using cDNA is via a GenomeWalker® kit, commercially available from Clontech (Palo Alto, CA), and described on page 27 of the 1997-1998 Clontech catalog.

Targeting Sequences: Non-Essential Genes

5 Central to the present invention is the fact that the targeting constructs contain "targeting" sequences (flanking, for example, the light generating protein-encoding sequence and promoter) derived from a single-copy, non-essential gene. These targeting sequences in the construct act via homologous recombination to replace at least a portion of the non-essential gene in the genome with the light-generating protein-encoding (*e.g.*,
10 luciferase-encoding) sequence operably linked to a promoter.

Non-limiting examples of targeting sequences for use in generating transgenic mice include sequences obtained from or derived from vitronectin, Fos B and galactin 3. A search of Mouse Knockout & Mutation Database (Genome Systems, Inc., St. Louis, MO) can be used to identify genes that have been knocked-out in mice where the
15 generated knockout mice displayed no obvious defects. The chromosomal locus for all these genes can be used to target promoter-(light generating protein, *e.g.*, luciferase) transgenes similar to what is described in Example 2. Single-copy, non-essential mouse genes identified in this manner include, but are not limited to, the following: Moesin (Msn), Doi Y., et al., J Biol Chem 1999, 274:2315-2321; Plasminogen activator
20 inhibitor, type II (Planh2) and Planh1, Dougherty K.M., Proc Natl Acad Sci USA 1999, 96:686-691; Protein tyrosine phosphatase, receptor type, B (Ptpnb), Elchebly et al. (1999) Science 283:1544-1548; Presenilin 1 (Psen1), Guo Q, et al. (1999) Proc Natl Acad Sci USA, 96:4125-4130; Protein kinase, mitogen-activated 9 (Prkm9) / SAPK/Erk/kinase 2 (Serk2), Kuan CY et al. (1999) Neuron 4:667-676; CD152 antigen
25 (Cd152) / CD86 antigen (Cd86) / CD80 antigen (Cd80), Mandelbrot DA, et al. (1999) J Exp Med, 189:435-440; Poly (ADP-ribose) polymerase (Adprp), Masutani M, et al. (1999), Proc Natl Acad Sci USA 96:2301-2304; Sodium channel, nonvoltage-gated 1 beta (Scnn1b), Pradervand S, et al. (1999) Proc Natl Acad Sci USA 96:1732-1737; Nuclear receptor coactivator 1 (Nco1), Qi C, et al. (1999) Proc Natl Acad Sci USA
30 96:1585-1590; Decay accelerating factor 1 (Daf1), Sun X, et al. (1999) Proc Natl Acad Sci USA 1999, 96:628-633; Necdin (Ndn), Tsai TF, et al. (1999) Nat Genet 22:15-16; Relaxin (Rln); Zhao L, et al. (1999) Endocrinology 140:445-453; Adenylyl cyclase 8

- (Adcy8), Abdel-Majid RM, et al. (1998) *Nat Genet* 19:289–291; Leukemia inhibitory factor (Lif), Bugga L, et al. (1998) *J Neurobiol* 36:509–524; Lectin, galactose binding, soluble 3 (Lgals3) and Lgals1, Calnot C, et al. (1998) *Dev Dyn* 211:306–313; Urokinase plasminogen activator receptor (Plaur) Carmeliet P, et al. (1998) *J Cell Biol* 140:233–
- 5 245; Nitric oxide synthase 1, neuronal (Nos1), Chao DS, et al. (1998) *J Neurochem* 71:784–789; Homeo box A7 (Hoxa7), Chen F, et al. (1998) *Mech Dev* 77:49–57; Myosin light chain, phosphorylatable, cardiac ventricles (MyIpc) Chen J, et al. (1998) *J Biol Chem* 273:1252–1256; Homeo box B7 (Hoxb7), Chen F, et al. (1998) *Mech Dev* 77:49–57; Nuclear factor of kappa light chain gene enhancer in B-cells inhibitor, alpha and beta (Nfkb1a and Nfkb1b), Cheng JD, et al. (1998) *J Exp Med* 186:1055–1062; Enolase
- 10 1, alpha non-neuron (Eno1), Couldrey C, et al. (1998) *Dev Dyn* 212:284–292; Xeroderma pigmentosum, complementation group A (Xpa), De Vries A, et al. (1998) *Exp Eye Res* 1998, 67:53–59; Von Willebrand factor homolog (Vwf), Denis C, et al. (1998) *Proc Natl Acad Sci USA* 95:9524–9529; Lysosomal acid lipase 1 (Lip1), Du H, et
- 15 al. (1998) *Hum Mol Genet* 7:1347–1354; UNC-5 homolog (*C. elegans*) 3 (Unc5h3), Eisenman LM, et al. (1998) *J Comp Neurol* 394:106–117; Protein phosphatase 1, regulatory (inhibitor) subunit 1B (Ppp1r1b), Fienberg AA, et al. (1998) *Science* 281:838–842; Myelin-associated glycoprotein (Mag) Fujita N, et al. (1998) *J Neurosci* 18:1970–1978; Paraoxonase 1 (Pon1), Furlong CE, et al. (1998) *Neurotoxicology*
- 20 19:645–650; Brain derived neurotrophic factor (Bdnf), Garek RR, et al. (1998) *Laryngoscope* 108:671–678; Neurotrophin 3 (Ntf3), Garek RR, et al. (1998) *Laryngoscope* 168:671–678; Myoglobin (Mb); Garry DJ, et al. (1998) *Nature* 395:905–908; Opioid receptor, mu (Oprm), Gaveriaux-Ruff C, et al. (1998) *Proc Natl Acad Sci USA* 95:6326–6330; Neuropeptide Y (Npy), Hollopeter G, et al. (1998) *Int J Obes Relat*
- 25 *Metab Disord* 22:506–512; Procollagen, type I, alpha 1 (Col1a1) Hormuzdi SG, et al. (1998) *Mol Cell Biol* 18:3368–3375; Centromere autoantigen B (Cenpb), Hudson DF, et al. (1998) *J Cell Biol* 141:309–319; Oculocerebrorenal syndrome of Lowe (ocrl), Janne PA, et al. (1998) *J Clin Invest* 101:2042–2053; arachidonate 12-lipoxygenase (Alox12) Johnson EN, et al. (1998) *Proc Natl Acad Sci USA* 95:3100–3105; H19 fetal liver
- 30 mRNA (H19), Jones BK, et al. (1998) *Genes Dev* 12:2200–2207; Hepatocyte nuclear factor 3 gamma (winged helix transcription factor) (Hnf3g), Kaestner KH, et al. (1998) *Mol Cell Biol* 18:4245–4251; Bone morphogenetic protein 2 (Bmp2) / Bone

- morphogenetic protein 7 (Bmp7) and Bmp5, Katagiri T, et al. (1998) *Dev Genet* 22:340–348; Intercellular adhesion molecule (Icam1), Ley K, et al. (1998) *Circ Res* 83:287–294; Glutamyl aminopeptidase (Enpep), Lin Q, et al. *J Immunol* 1998, 160:4681–4687; Prion protein (Prnp), Lipp HP, et al. *Behav Brain Res* 1998, 95:47–54; RAB3A, member RAS
- 5 oncogene family (Rab3a), Lonart G, et al. *Neuron* 1998, 21:1141–1150; Potassium voltage gated channel, shaker related subfamily, member 4 (Kcna4), London B, et al., *J Physiol (Lond)* 1998, 509:171–182; Apurinic/apurimidinic endonuclease (Apex), Ludwig DL, et al. *Mutat Res* 1998, 409:17–29; T-cell receptor gamma, variable 5 (Tcrg-V5), Mallick-Wood CA, et al. *Science* 1998, 279:1729–1733; Nuclear, factor, erythroid
- 10 derived 2, like 2 (Nfe2l2), Martin F, et al. *Blood* 1998, 91:3459–3466; Interleukin 13 (Il13), McKenzie GJ, et al. *Curr Biol* 1998, 8:339–340; Sorbitol dehydrogenase 1 (Sdh1), Ng T, et al. *Diabetes* 1998, 47:961–966; Guanine nucleotide binding protein, alpha 11 (Gna11), Offermanns S, et al. *EMBO J* 1998, 17:4304–4312; Estrogen receptor alpha (Estra), Ogawa S, et al. *Endocrinology* 1998, 139:5070–5081; Integrin beta 2 (Itgb2),
- 15 Intracellular adhesion molecule (Icam1) and CD34 antigen, Oliveira-dos-Santos AJ, et al. *Eur J Immunol* 1998, 28:2882–2892; Angiotensin receptor 1b (Agt1b), Oliverio MI, et al. *Proc Natl Acad Sci USA* 1998, 95:15496–15501; Complement factor B (factor B), Pekna M, et al. *Scand J Immunol* 1998, 47:375–380; Centromere autoantigen B (Cenpb), Perez-Castro AV, et al. *Dev Biol* 1998, 201:135–143; Procollagen, type V, alpha 2
- 20 (Col5a2) / Fibrillin 1 (Fbn1), Phelps RG, et al. *Mol Med* 1998, 4:356–360; Plasminogen activator inhibitor, type 1 (Planh1), Pinsky DJ, et al. *J Clin Invest* 1998, 102:919–928, Carmeliet P, et al. *J Clin Invest* 1993, 92:2756–2760; Placentae and embryos oncofetal gene (Pem), Pitman JL, et al. *Dev Biol* 1998, 202:196–214; Postmeiotic segregation increased 1 (*S. cerevisiae*) (Pms1), Prolla TA, et al. *Nat Genet* 1998, 18:276–278; Prion
- 25 protein, structural locus (Prn-p), Prusiner SB, et al. *Proc Natl Acad Sci USA* 1998, 90:10608–10612, Lledo P-M, et al. *Proc Natl Acad Sci U S A* 1996, 93:2403–2407, Sailer A, et al. *Cell* 1994, 77:967–968, Weissmann C, et al. *Philos Trans R Soc Lond [Biol]* 1994, 343:431–433, Bueler H, et al. *Cell* 1993, 73:1337–1347, Weissmann C, et al. *Intervirology* 1993, 35:164–175; NAD (P)H:quinone oxidoreductase, Radjendirane V,
- 30 et al. *J Biol Chem* 1998, 273:7382–7389; Alpha tropomyosin (Tpm1), Rethinasamy P, et al. *Circ Res* 1998, 82:116–123; Goosecoid and Goosecoid-like (Gsc1), Wakamiya M, et al. *Hum Mol Genet* 1998, 7:1835–1840, Saint-Pore B, et al. *Hum Mol Genet* 1998,

- 7:1841–1849; Schlafen 1 (Slfn1); Schwarz DA, et al. *Immunity* 1998, 9:657–668; Nuclear factor, erythroid derived 2, ubiquitous (Mafk), Shavit JA, et al. *Genes Dev* 1998, 12:2164–2174; Microphthalmia-associated transcription factor (Mitf), Smith SB, et al. *Exp Eye Res* 1998, 66:403–410; Bone morphogenetic protein 6 (Bmp6), Solloway MJ, et al. *Dev Genet* 1998, 22:321–339; Phosphatidylinositol glycan, class A (Piga), Takahama Y, et al. *Eur J Immunol* 1998, 28:2159–2166; Paired-related homeobox 2 (Prx2), Berge D, et al. *Development* 1998, 125:3831–3842; Prostaglandin E receptor EP1 subtype (Ptgerep1), Ushikubi F, et al. *Nature* 1998, 395:281–284; Immunoglobulin kappa chain complex (Igk), van der Stoep N, et al. *Immunity* 1998, 8:743–750; Adenine phosphoribosyl transferase (Aprt), Van Sloun PP, et al. *Nucleic Acids Res* 1998, 26:4888–4894; Microtubule associated protein 4 (Mtap4), Voss AK, et al. *Dev Dyn* 1998, 212:258–266; ; 3-hydroxy-3-methylglutaryl-coenzyme A lyase (Hmgcl), Wang SP, et al. *Hum Mol Genet* 1998, 7:2057–2062; Fibroblast growth factor receptor 4 (Fgfr4), Weinstein M, et al. *Development* 1998, 125:3615–3623; Hepsin (Hpn), Wu Q, et al. *J Clin Invest* 1998, 101:321–326; Small inducible cytokine A11 (Scya11), Yang T, et al. *Blood* 1998, 92:3912–3923; Small nuclear ribonucleoprotein N (Snrpn), Yang T, et al. *Nat Genet* 1998, 19:25–31; DNA fragmentation factor, alpha subunit (Dffa), Zhang J, et al. *Proc Natl Acad Sci USA* 1998, 95:12480–12485; Early growth response 1 (Egr1), Zheng D, et al. *Neuroscience* 1998, 83:251–258; Early growth response 1 (Egr1) /
- Hormone receptor (Hmr), Zheng D, et al. *Neuroscience* 1998, 83:251–258; Hemochromatosis (Hfe), Zhou XY, et al. *Proc Natl Acad Sci USA* 1998, 95:2492–2497; Alpha tropomyosin (Tpm1), Blanchard EM, et al. *Circ Res* 1997, 81:1005–1010; tRNA phosphoserine (Trsp), Bosl MR, et al. *Proc Natl Acad Sci U S A* 1997, 94:5531–5534; Angiotensin receptor 1b (Agtr1b), Chen X, et al. *Am J Physiol* 1997, 272:F299–F304;
- Xeroderma pigmentosum, complementation group C (Xpc), Cheo DL, et al. *Mut Res* 1997, 374:1–9; B cell leukemia/lymphoma 6 (Bcl6), Dent AL, et al. *Science* 1997, 276:589–592; Fumarylacetoacetate hydrolase (Fah) / 4-hydroxyphenylpyruvic acid dioxygenase (Hpd), Endo F, et al. *J Biol Chem* 1997, 272:24426–24432; N-methylpurine-DNA glycosylase (Mpg), Engelward BP, et al. *Proc Natl Acad Sci USA* 1997, 94:13087–13092; Interleukin 1 receptor, type 1 (Il1r1), Glaccum MB, et al. *J Immunol* 1997, 159:3364–3371; N-methylpurine-DNA glycosylase (Mpg), Hang B, et al. *Proc Natl Acad Sci USA* 1997, 94:12869–12874; Gamma-aminobutyric acid (GABA-

- A) receptor, subunit alpha 6 (Gabra6), Homanics GE, et al. *Mol Pharmacol* 1997, 51:588–596; Superoxide dismutase 2, mitochondrial (Sod2), Huang TT, et al. *Arch Biochem Biophys* 1997, 344:424–432; Interleukin 11 receptor, alpha chain 1 (Il11ra1), Nandurjar HH, et al. *Blood* 1997, 90:2148–2159; Alkaline phosphatase 5 (Akp5),
- 5 Narisawa S, et al. *Dev Dyn* 1997, 208:432–446; GATA-binding protein 4 (Gata4), Narita N, et al. *Development* 1997, 124:3755–3764; Lymphocyte protein tyrosine kinase (Lck) / Fyn protooncogene (Fyn); Page ST, et al. *Eur J Immunol* 1997, 27:554–562; P glycoprotein 3 (Pgy3), Schinkel AH, et al. *Proc Natl Acad Sci U S A* 1997, 94:4028–4033; P glycoprotein 1 (Pgy1) / P glycoprotein 3 (Pgy3), Schinkel AH, et al., *Proc Natl*
- 10 *Acad Sci U S A* 1997, 94:4028–4033; Creatine kinase, mitochondrial 1, ubiquitous (Ckmt1), Steeghs K, et al. *J Neurosci Methods* 1997, 71:29–41; T cell receptor gamma, variable 4 (Tcrg-V4), Sunaga S, et al. *J Immunol* 1997, 158:4223–4228; Ia-associated invariant chain (p31 form) (Ii), Takaesu NT, et al. *J Immunol* 1997, 158:187–199; Solute carrier family 18 (vesicular monoamine), member 2 (Slc18a2), Takahashi N, et al. *Proc*
- 15 *Natl Acad Sci USA* 1997, 94:9938–9943; Matrix metalloproteinase 7 (Mmp7), Wilson CL, et al. *Proc Natl Acad Sci U S A* 1997, 94:1402–1407; Formin (Fmn), Wynshaw-Boris A, et al. *Mol Med* 1997, 3:372–384; Synaptophysin (Syp), Arrandale JM, et al. *J Biol Chem* 1996, 271:21353–21358; Transformation related protein 53 (Trp53), Boehme SA, et al. *J Immunol* 1996, 156:4075–4078; Neuronal nitric oxide synthase (Nos1),
- 20 Burnett AL, et al. *Mol Medicine* 1996, 2:288–296; Eph receptor A2 (Epha2), Chen J, et al. *Oncogene* 1996, 12:979–988; Urokinase plasminogen activator receptor (Plaur); Dewerchin M, et al. *J Clin Invest* 1996, 97:870–878; Growth differentiation factor 9 (Gdf9), Dong J, et al. *Nature* 1996, 383:531–535; Externally regulated phosphatase (Ptpn16), Dorfman K, et al. *Oncogene* 1996, 13:925–931; Tenascin C (Tnc), Forsberg E,
- 25 et al. *Proc Natl Acad Sci U S A* 1996, 93:6594–6599; Integrin alpha 1 (Itga1), Gardner H, et al. *Dev Biol* 1996, 175:301–313; FBJ osteosarcoma oncogene B (Fosb), Gruda MC, et al. *Oncogene* 1996, 12:2177–2185; Breast cancer 1 (Brca1), Hakem R, et al. *Cell* 1996, 85:1009–1023; Megakaryocyte-associated tyrosine kinase (Matk), Hamaguchi I, et al. *Biochem Biophys Res Commun* 1996, 224:172–179; Apolipoprotein B editing
- 30 complex 1 (Apobec1), Hirano K-I, et al. *J Biol Chem* 1996, 271:9887–9890; Carboxyl ester lipase, Howles PN, et al. *J. Biol. Chem.* 1996, 271:7196–7202; Nuclear factor, erythroid derived 2, ubiquitous (Nfe2u), Kotkow KJ, Orkin SH. *Proc Natl Acad Sci U S*

- A 1996, 93:3514–3518; Retinoid X receptor gamma (Rxrg), Krezel W, et al. *Proc Natl Acad Sci U S A* 1996, 93:9010–9014; Early growth response 1 (Egr1), Lee SL, et al. *Mol Cell Biol* 1996, 16:4566–4572; Adrenergic receptor, alpha 2b (Adra2b), Link RE, et al. *Science* 1996, 273:803–805; Angiotensin receptor 1a (Agtr1a), Matsusaka T, et al. *J Clin Invest* 1996, 98:1867–1877; Interleukin 3 receptor, beta chain 1 (Il3rb2), Nicola NA, et al. *Blood* 1996, 87:2665–2674; Transformation related protein 53 (Trp53), Ohashi M, et al. *Jpn J Cancer Res* 1996, 87:696–701; Leukemia-associated gene (Lag), Schubart U K, et al. *J Biol Chem* 1996, 271:14062–14066; Glutathione peroxidase 1 (Gpx1), Spector A, et al. *Exp Eye Res* 1996, 62:521–540; Arachidonate 12-lipoxygenase, leukocyte (Alox12l), Sun D, Funk CD. *J Biol Chem* 1996, 271:24055–24062; Complement component 3 (C3), Sylvestre D, et al. *J Exp Med* 1996, 184:2385–2392; CD30 antigen (Cd30), Texido G, et al. *Eur J Immunol* 1996, 26:1966–1969; Fc receptor, IgE, low affinity II, alpha polypeptide (Fcer2a), Immunoglobulin heavy chain 5 (delta-like heavy chain) (Igh-5), Interleukin 4 (IL4) and Terminal-deoxynucleotidyl transferase, Texido G, et al. *Eur J Immunol* 1996, 26:1966–1969; Apolipoprotein A-II (Apoa2), Weng W, Breslow JL. *Proc Natl Acad Sci U S A* 1996, 93:14788–14794; Amyloid beta (A4) precursor protein (App), Zheng H, et al. *Ann N Y Acad Sci* 1996, 777:421–426; cAMP responsive element binding protein 1 (Creb1), Blendy JA, et al. *Brain Res* 1995, 681:8–14; Bradykinin receptor, beta 2 (Bdkrb2), Borkowski JA, et al. *J Biol Chem* 1995, 270:13706–13710; Growth factor response protein (Gfrp), Crawford PA, et al. *Mol Cell Biol* 1995, 15:4331–4336; Ciliary neurotrophic factor (Cntf), de Chiara TM, et al. *Cell* 1995, 83:313–322; Cyclin dependent kinase inhibitor 1A (P21) (Cdkn1a); Deng C, et al. *Cell* 1995, 82:675–684; Granzyme A (Gzma), Ebnet K, et al. *EMBO J* 1995, 14:4230–4239; Very low density lipoprotein receptor (Vldlr), Frykman PK, et al. *Proc Natl Acad Sci U S A* 1995, 92:8453–8457; Apolipoprotein E (ApoE) / Apolipoprotein A-I (Apoa1), Goodrum JF, et al. *J Neurobiol* 1995, 64:408–416; Nitric oxide synthase 1, neuronal (Nos1), Ichinose F, et al. *Anesthesiology* 1995, 83:101–108; Nitric oxide synthase 2, inducible, macrophage (Nos2), Laubach VE, et al. *Proc Natl Acad Sci U S A* 1995, 92:10688–10692; Peroxisome proliferator activated receptor alpha (Ppara), Lee SS-T, et al. *Mol Cell Biol* 1995, 15:3012–3022; Growth factor response protein (Gfrp), Lee SL, et al. *Science* 1995, 269:532–535; H19 fetal liver mRNA (H19) / Insulin-like growth factor 2 (Igf2), Leighton PA, et al. *Nature* 1995, 375:34–39; Retinoic

- acid receptor beta (RAR-beta), Luo J, et al. *Mech Dev* 1995, 53:61-71; Metallothionein 1 (Mt1) / Metallothionein 2 (Mt2), Philcox JC, et al. *Biochem J* 1995, 308:543-546; Heme oxygenase (decycling) 2 (Hmox2), Poss KD, et al. *Neuron* 1995, 15:867-873; H1-0 histone (H1fv), Sirotkin AM, et al. *Proc Natl Acad Sci U S A* 1995, 92:6434-6438;
- 5 Creatine kinase, mitochondrial 1, ubiquitous (Ckmt1), Steeghs K, et al. *Biochim Biophys Acta* 1995, 1230:130-138; Tenascin C (Tnc), Steindler DA, et al. *J Neurosci* 1995, 15:1971-1983; Ia-associated invariant chain (Ii), Takaesu NT, et al. *Immunity* 1995, 3:385-396; Neuroblastoma ras oncogene (Nras), Umanoff H, et al. *Proc Natl Acad Sci USA* 1995, 92:1709-1713; Receptor-associated protein of the synapse, 43 kDa (Rapsn),
- 10 Willnow TE, et al. *Proc Natl Acad Sci USA* 1995, 92:4537-4541; Vitronectin (Vtn), Zheng X, et al. *Proc Natl Acad Sci U S A* 1995, 92:12426-12430; Preproacrosin (Acr), Baba T, et al. *J Biol Chem* 1994, 269:31845-31849; Vimentin (Vim), Colucci GE, et al. *Cell* 1994, 79:679-694; Tumor necrosis factor receptor superfamily, member 1b (Tnfrsf1b), Erickson SL, et al. *Nature* 1994, 372:560-563; Cellular retinoic acid binding
- 15 protein I (Crabp1), Gorry P, et al. *Proc Natl Acad Sci USA* 1994, 91:9032-9036; cAMP responsive element binding protein 1 (Creb1), Hummler F, et al. *Proc Natl Acad Sci USA* 1994, 91:5647-5651; Pore forming protein (Pfp), Kagi D, et al. *Nature* 1994, 369:31-37; Src-related kinase lacking C-terminal regulatory tyrosine and N-terminal myristylation sit (Srms), Kohmura N, et al. *Mol Cell Biol* 1994, 14:6915-6925; CD3
- 20 polypeptide zeta (Cd3z) / Solute carrier family 22, member 1 (Slc22a1), Koyasu S, et al. *EMBO J* 1994, 13:784-797; Pore forming protein (Pfp), Lowin B, et al. *Proc Natl Acad Sci USA* 1994, 91:11571-11575; Retinoic acid receptor, beta (Rarb), Retinoic acid receptor beta2 (RARbeta2), Mendelsohn C, et al. *Dev Biol* 1994, 166:246-258; Transthyretin (Ttr), Palha JA, et al. *J Biol Chem* 1994, 269:33135-33139; Procollagen,
- 25 type X, alpha 1 (Col10a1), Rosati R, et al. *Nature Genet* 1994, 8:129-135; P glycoprotein 3 (Pgy3), Schinkel AH, et al. *Cell* 1994, 77:491-502; Yamaguchi sarcoma viral (v-yes) oncogene homolog (Yes), Stein PL, et al. *Genes Dev* 1994, 8:1999-2007; Fc receptor, IgE, high affinity II, alpha polypeptide (Fcer2a), Stief A, et al. *J Immunol* 1994, 152:3378-3390; Pore forming protein (Pfp), Walsh CM, et al. *Proc Natl Acad Sci*
- 30 *USA* 1994, 91:10854-10858; CD2 antigen (Cd2), Evans CF, et al. *J Immunol* 1993, 151:6259-6264; Mannose-6-phosphate receptor, cation dependent (M6pr), Koster A, et al. *EMBO J* 1993, 12:5219-5223; Retinoic acid receptor, alpha (Rara), Li E, et al. *Proc*

- Natl Acad Sci USA 1993, 90:1590–1594; Lufkin T, et al. Proc Natl Acad Sci USA 1993, 90:7225–7229; Retinoic acid receptor, gamma (Rarg), Lohnes D, et al. Cell 1993, 73:643–658; Tumor necrosis factor receptor 1 (TNF-R-1) (Tnfr1), Pfeffer K, et al. Cell 1993, 73:457–467; Lectin, galactose binding, soluble 1 (Lgals1), Poirier F, Robertson EJ.
- 5 Development 1993, 119:1229–1236; Synapsin I (Syn1), Rosahl TW, et al. Cell 1993, 75:661–670; Tumor necrosis factor receptor 1 (Tnfr1), Rothe J, et al. Nature 1993, 364:798–802; Beta-2 microglobulin (B2m), Correa I, et al. Proc Natl Acad Sci USA 1992, 89:653–657; CD2 antigen (Cd2), Killeen N, et al. EMBO J 1992, 11:4329–4336; Apolipoprotein E (ApoE), Piedrahita JA, et al. Proc Natl Acad Sci USA 1992, 89:4471–
- 10 4475; Myogenic differentiation 1 (Myod1), Rudnicki MA, et al. Cell 1992, 71:383–390; Tenascin C (Tnc), Saga Y, et al. Genes Dev 1992, 6:1821–1831; Beta-2 microglobulin (B2m), Sanjuan N, et al. J Virol 1992, 66:4587–4590; Neuroblastoma myc-related oncogene 1 (Nmyc1), Stanton BR, et al. Genes Dev 1992, 6:2235–2247; and Hemoglobin alpha chain complex (Hba), Popp RA, et al. Genetics 1983, 105:157–167.
- 15 Some preferred single-copy, non-essential genes with no phenotypes of the present invention include, but are not limited to, the following: Moesin (Msn), Doi Y., et al., J Biol Chem 1999, 274:2315–2321; Plasminogen activator inhibitor, type II (Planh2) and Planh1, Dougherty K.M., Proc Natl Acad Sci USA 1999, 96:686–691; Nuclear receptor coactivator 1 (Nco1), Qi C, et al. (1999) Proc Natl Acad Sci USA 96:1585–
- 20 1590; Nuclear factor of kappa light chain gene enhancer in B-cells inhibitor, alpha and beta (Nfkb1a and Nfkb1b), Cheng JD, et al. (1998) J Exp Med 6:1055–1062; H19 fetal liver mRNA (H19), Jones BK, et al. (1998) Genes Dev 12:2200–2207; Prion protein (Prnp), Lipp HP, et al. Behav Brain Res 1998, 95:47–54; Centromere autoantigen B (Cenpb), Perez-Castro AV, et al. Dev Biol 1998, 201:135–143; Placentae and embryos
- 25 oncofetal gene (Pem), Pitman JL, et al. Dev Biol 1998, 202:196–214; Externally regulated phosphatase (Ptpn16), Dorfman K, et al. Oncogene 1996, 13:925–931; Transformation related protein 53 (Trp53), Ohashi M, et al. Jpn J Cancer Res 1996, 87:696–701; H1-O histone (H1fv), Sirotkin AM, et al. Proc Natl Acad Sci U S A 1995, 92:6434–6438; Creatine kinase, mitochondrial 1, ubiquitous (Ckmt1), Steeghs K, et al.
- 30 Biochim Biophys Acta 1995, 1230:130–138; Neuroblastoma ras oncogene (Nras), Umanoff H, et al. Proc Natl Acad Sci USA 1995, 92:1709–1713; Vitronectin (Vtn), Zheng X, et al. Proc Natl Acad Sci U S A 1995, 92:12426–12430; Vimentin (Vim),

Colucci GE, et al. Cell 1994, 79:679–694; Cellular retinoic acid binding protein I (Crabp1), Gorry P, et al. Proc Natl Acad Sci USA 1994, 91:9032–9036; Retinoic acid receptor beta2 (RARbeta2), Mendelsohn C, et al. Dev Biol 1994, 166:246–258; Retinoic acid receptor, alpha (Rara), Li E, et al. Proc Natl Acad Sci USA 1993, 90:1590–1594,
5 Lufkin T, et al. Proc Natl Acad Sci USA 1993, 90:7225–7229; Lectin, galactose binding, soluble 1 (Lgals1), Poirier F, Robertson EJ. Development 1993, 119:1229–1236; Myogenic differentiation 1 (Myod1), Rudnicki MA, et al. Cell 1992, 71:383–390; and Tenascin C (Tnc), Saga Y, et al. Genes Dev 1992, 6:1821–1831.

In view of the guidance of the present specification, one of ordinary skill in the
10 art can select similar, suitable, single-copy, non-essential genes in mice and other cell types/organisms.

Assembly of Targeting Cassettes

The targeting cassettes described herein can be constructed utilizing
15 methodologies known in the art of molecular biology (see, for example, Ausubel or Maniatis) in view of the teachings of the specification. As described above, the targeting constructs are assembled by inserting, into a suitable vector backbone, polynucleotides encoding a reporter, such as a light-generating protein, *e.g.*, a luciferase gene, operably linked to a promoter of interest; a sequence encoding a positive selection marker; and,
20 optionally a sequence encoding a negative selection marker. In addition, the targeting cassette contains insertion sites such that sequences targeting a single-copy, non-essential gene can be readily inserted to flank the sequence encoding positive selection marker and luciferase-encoding sequence.

A preferred method of obtaining polynucleotides, suitable regulatory sequences
25 (*e.g.*, promoters) is PCR. General procedures for PCR as taught in MacPherson et al., PCR: A PRACTICAL APPROACH, (IRL Press at Oxford University Press, (1991)). PCR conditions for each application reaction may be empirically determined. A number of parameters influence the success of a reaction. Among these parameters are annealing temperature and time, extension time, Mg²⁺ and ATP concentration, pH, and the relative
30 concentration of primers, templates and deoxyribonucleotides. Exemplary primers are described below in the Examples. After amplification, the resulting fragments can be

detected by agarose gel electrophoresis followed by visualization with ethidium bromide staining and ultraviolet illumination.

In one embodiment, PCR can be used to amplify fragments from genomic libraries. Many genomic libraries are commercially available. Alternatively, libraries
5 can be produced by any method known in the art. Preferably, the organism(s) from which the DNA is has no discernible disease or phenotypic effects. This isolated DNA may be obtained from any cell source or body fluid (*e.g.*, ES cells, liver, kidney, blood cells, buccal cells, cerviovaginal cells, epithelial cells from urine, fetal cells, or any cells present in tissue obtained by biopsy, urine, blood, cerebrospinal fluid (CSF), and tissue
10 exudates at the site of infection or inflammation). DNA is extracted from the cells or body fluid using known methods of cell lysis and DNA purification. The purified DNA is then introduced into a suitable expression system, for example a lambda phage.

Another method for obtaining polynucleotides, for example, short, random nucleotide sequences, is by enzymatic digestion. As described below in the Examples,
15 short DNA sequences generated by digestion of DNA from vectors carrying genes encoding luciferase (yellow green or red).

Polynucleotides are inserted into vector genomes using methods known in the art. For example, insert and vector DNA can be contacted, under suitable conditions, with a restriction enzyme to create complementary or blunt ends on each molecule that can pair
20 with each other and be joined with a ligase. Alternatively, synthetic nucleic acid linkers can be ligated to the termini of a polynucleotide. These synthetic linkers can contain nucleic acid sequences that correspond to a particular restriction site in the vector DNA. Other means are known and, in view of the teachings herein, can be used.

25 The final constructs can be used immediately (*e.g.*, for introduction into ES cells), or stored frozen (*e.g.*, at -20°C) until use. Preferably, the constructs are linearized prior to use, for example by digestion with suitable restriction endonucleases.

Transgenic Animals

The targeting constructs containing the light generating protein coding sequences (e.g., luciferase genes) are introduced into a pluripotent cell (e.g., ES cell, Robertson, E. J., In: Current Communications in Molecular Biology, Capecchi, M. R. (ed.), Cold Spring Harbor Press, Cold Spring Harbor, N.Y. (1989), pp. 39-44). Suitable ES cells may be derived or isolated from any species or from any strain of a particular species. Although not required, the pluripotent cells are typically derived from the same species as the intended recipient. ES cells may be obtained from commercial sources, from International Depositories (e.g., the ATCC) or, alternatively, may be obtained as described in Robertson, E. J., *supra*. Examples of clonally-derived ES cells lines include 129/SVJ ES cells, RW-4 and C57BL/6 ES cells (Genome Systems, Inc.).

ES cells are cultured under suitable conditions, for example, as described in Ausubel et al., section 9.16, *supra*. Preferably, ES cells are cultured on stromal cells (such as STO cells (especially SNC4 STO cells) and/or primary embryonic fibroblast cells) as described by E. J. Robertson, *supra*, pp 71-112. Culture media preferably includes leukocyte inhibitory factor ("lif") (Gough, N. M. et al., Reprod. Fertil. Dev. 1:281-288 (1989); Yamamori, Y. et al., Science 246:1412-1416 (1989), which appears to help keep the ES cells from differentiating in culture. Stromal cells transformed with the gene encoding lif can also be used.

The targeting constructs are introduced into the ES cells by any method which will permit the introduced molecule to undergo recombination at its regions of homology, for example, micro-injection, calcium phosphate transformation, or electroporation (Toneguzzo, F. et al., Nucleic Acids Res. 16:5515-5532 (1988); Quillet, A. et al., J. Immunol. 141:17-20 (1988); Machy, P. et al., Proc. Natl. Acad. Sci. (U.S.A.) 85:8027-8031 (1988)). The construct to be inserted into the ES cell must first be in the linear form. Thus, if the knockout construct has been inserted into a vector as described above, linearization is accomplished by digesting the DNA with a suitable restriction endonuclease selected to cut only within the vector sequence and not within the knockout construct sequence. If the ES cells are to be electroporated to insert the construct, the ES cells and construct DNA are exposed to an electric pulse using an electroporation machine and following the manufacturer's guidelines for use. After electroporation, the ES cells are typically allowed to recover under suitable incubation conditions. The cells

are then cultured under conventional conditions, as are known in the art, and screened for the presence of the construct.

Screening and selection of those cells into which the targeting construct has been integrated can be achieved using the positive selection marker and/or the negative
5 selection marker in the construct. In preferred embodiments, the construct contains both positive and negative selection markers. In one aspect, methods which rely on expression of the selection marker are used, for example, by adding the appropriate substrate to select only those cells which express the product of the positive selection marker or to eliminate those cells expressing the negative selection marker. For
10 example, where the positive selection marker encodes neomycin resistance, G418 is added to the transformed ES cell culture media at increasing dosages. Similarly, where the negative selection marker is used, a suitable substrate (*e.g.*, gancyclovir if the negative selection marker encodes HSV-TK) is added to the cell culture. Either before or after selection using the appropriate substrate, the presence of the positive and/or
15 negative selection markers in a recipient cell can also be determined by others methods, for example, hybridization, detection of radiolabelled nucleotides, PCR and the like. In preferred embodiments, cells having integrated targeting constructs are first selected by adding the appropriate substrate for the positive and/or negative selection markers. Cells that survive the selection process are then screened by other methods, such as PCR or
20 Southern blotting, for the presence of integrated sequences.

After suitable ES cells containing the construct in the proper location have been identified, the cells can be inserted into an embryo, preferably a blastocyst. The blastocysts are obtained by perfusing the uterus of pregnant females. In one embodiment, the blastocysts are obtained from, for example, the FVB/N strain of mice and the ES cells
25 are obtained from, for example, the C57BL/6 strain of mice. Suitable methods for accomplishing this are known to the skilled artisan, and are set forth by, *e.g.*, Bradley et al., (1992) *Biotechnology*, 10:534-539. Insertion into the embryo may be accomplished in a variety of ways known to the skilled artisan, however a preferred method is by microinjection. For microinjection, about 10-30 ES cells are collected into a micropipet
30 and injected into embryos that are at the proper stage of development to permit integration of the foreign ES cell containing the construct into the developing embryo.

The suitable stage of development for the embryo used for insertion of ES cells is species dependent, in mice it is about 3.5 days.

While any embryo of the right stage of development is suitable for use, it is preferred that blastocysts are used. In addition, preferred blastocysts are male and, 5 furthermore, preferably have genes encoding a coat color that is different from that encoded by the genes ES cells. In this way, the offspring can be screened easily for the presence of the knockout construct by looking for mosaic coat color (indicating that the ES cell was incorporated into the developing embryo). Thus, for example, if the ES cell line carries the genes for black fur, the blastocyst selected will carry genes for white or 10 brown fur.

After the ES cell has been introduced into the blastocyst, the blastocyst is typically implanted into the uterus of a pseudopregnant foster mother for gestation. Pseudopregnant females are prepared by mating with vasectomized males of the same species and successful implantation usually must occur within about 2-3 days of mating.

15 Offspring are screened initially for mosaic coat color where the coat color selection strategy has been employed. Southern blots and/or PCR may also be used to determine the presence of the sequences of interest. Mosaic (chimeric) offspring are then bred to each other to generate homozygous animals. Homozygotes and heterozygotes may be identified by Southern blotting of equivalent amounts of genomic DNA from 20 mice that are the product of this cross, as well as mice that are known heterozygotes and wild type mice. Alternatively, Northern blots can be used to probe the mRNA to identify the presence or absence of transcripts encoding either the replaced gene, the light generating protein coding sequence (e.g., luciferase gene), or both. In addition, Western blots can be used to assess the level of expression of the luciferase protein with an 25 antibody against the luciferase gene product. Finally, in situ analysis (such as fixing the cells and labeling with antibody) and/or FACS (fluorescence activated cell sorting) analysis of various cells from the offspring can be conducted using suitable (e.g., anti-luciferase) antibodies to look for the presence or absence of the targeting construct.

In one embodiment of the present invention, the animals are from the C57BL/6 30 mouse strain. This strain develops a variety of tumors and has been used to develop a number of tumor cells lines, for example, B16 melanoma cells (including, B16F10, B16D5, and B16F1), Lewis lung carcinoma cells (including, LLC, LLC-h59), T241

mouse fibrosarcoma cells, RM-1 and pTC2 mouse prostate cancer cells, and MCA207 mouse sarcoma cells. These cell lines have been extensively used for *in vivo* tumor biology studies after injection into C57BL/6 mice. The generated targeted transgenic mice in the Examples are in C57BL/6 genetic background and these animals are suitable
5 for injection or implantation of such tumor cells, as well as other tumor cells described in literature that are immunocompatent for C57BL/6 mice. Thus, the transgenic animals can then be used, for example, to monitor, *in vivo*, tumor progression (*e.g.*, growth) and the efficacy of therapies on tumor regression. For example, where the transgenic animal is tumor-susceptible, it is monitored for expression of a reporter, *e.g.*, luciferase, which is
10 indicative of tumorigenesis and/or angiogenesis. The monitoring of expression of luciferase reporter expression cassettes using non-invasive whole animal imaging has been described (Contag, C. et al, U.S. Patent No. 5,650,135, July 22, 1997; Contag, P., et al, *Nature Medicine* 4(2):245-247, 1998; Contag, C., et al, *OSA TOPS on Biomedical Optical Spectroscopy and Diagnostics* 3:220-224, 1996; Contag, C.H., et al,
15 *Photochemistry and Photobiology* 66(4):523-531, 1997; Contag, C.H., et al, *Molecular Microbiology* 18(4):593-603, 1995). Such imaging typically uses at least one photo detector device element, for example, a charge-coupled device (CCD) camera.

The transgenic animals described herein can also be used to determine the effect of an analyte (*e.g.*, therapy), for example on tumor progression where the promoter
20 induces light generating protein (*e.g.*, luciferase) expression when a tumor develops. Methods of administration of the analyte include, but are not limited to, injection (subcutaneously, epidermally, intradermally), intramucosal (such as nasal, rectal and vaginal), intraperitoneal, intravenous, oral or intramuscular. Other modes of administration include oral and pulmonary administration, suppositories, and transdermal
25 applications. Dosage treatment may be a single dose schedule or a multiple dose schedule. For example, the analyte of interest can be administered over a range of concentration to determine a dose/response curve. The analyte may be administered to a series of test animals or to a single test animal (given that response to the analyte can be cleared from the transgenic animal).

30 The following examples are intended only to illustrate the present invention and should in no way be construed as limiting the subject invention.

EXAMPLES

Example 1 Generating the Targeting Cassette and Vector

5

A. Creation of the Backbone Vector

pTK53: The 0.5 kb mouse phosphoglycerate kinase 1 promoter was amplified with PGK primers (PGKF, SEQ ID NO:1:

ATCGAATTCTACCGGGTAGGGGAGGCGCTTT; PGKR, SEQ ID NO:2:

10 GGCTGCAGGTCGAAAGGCCCGGAGATGAGG) using mouse genomic DNA (Genome Systems, Inc., St. Louis, MO) as template. This fragment was then double digested with EcoRI and PstI and cloned into the pKS vector (Stratagene, La Jolla, California) which was linearized with the same enzymes. The neomycin gene was amplified with NeoF (SEQ ID NO:3:

15 ACCTGCAGCCAATATGGGATCGGCCATTGAAC) and NeoR (SEQ ID NO:4: GGATCCGCGGCCCGCCCCAGCTGGTTCTTCCGCCTC) primers using pNTKV1907 (Stratagene) as a template. The 1.1 kb PCR fragment was double digested with PstI and BamHI and cloned into the pKS-PGK vector which was linearized with the same enzymes. This pKS-PGK-Neo vector was used to clone thymidine kinase gene as
20 follows. Primers TKF (SEQ ID NO: 5:

GGATCCTCTAGAGTCGAGCAGTGTGGTTTT) and TKR (SEQ ID NO:6:

GAGCTCCCGTAGTCAGGTTTAGTTCGTCCG) were used to amplify the TK gene from pNTKV1907 (Stratagene). The amplified 2kb fragment was then digested with

BamHI and SacI and cloned into pKS-PGK-Neo vector that was linearized with the same

25 enzymes. This constructed vector was designated as pTK. A synthetic linker F5R5 was made after annealing of two primers (forward primer, F5R51, SEQ ID NO: 7:

GTACATTTAAATCCTGCAGG; reverse primer, F5R52, SEQ ID NO:8:

AGCTCCTGCAGGATTTAAAT). This linker was inserted between Asp718I and

HindIII sites of pTK and the new construct was designated pTK5. A second synthetic

30 linker F3R3 was made by annealing of two primers (forward primer, F3R31, SEQ ID NO:9:

GGCCCGGGCTTAATTAATGCATCATATGGTACCGTTTAAACGCGGCCGCAAG

CTTGTCGACGGCGCGCCGGCCGGCC; reverse primer, F3R32, SEQ ID NO:10:

GATCGGCCGGCCGGCGCGCCGTCGACAAGCTTGCGGCCGCGTTTAAACGGTACCATATGATGCATTAATTAAGCCCG).). This linker was inserted between NotI and BamHI sites of pTK and the new construct was designated pTK53. Schematics of the vectors are shown in Figure 1.

5

B. Introduction of Luciferase

pTK-LucYG and pTK-LucR: The yellow green luciferase gene was isolated from pGL3 vector (Promega) as a HindIII-SalI fragment and was cloned into pGK53 that was linearized with the same enzymes. The new construct was designated pTK-LucYG (8931 bp), shown in Figure 2.

10

The red luciferase gene was isolated from pGL3-red vector (Dr. Christopher Contag, Stanford University, Stanford, Calif.) as a HindIII-SalI fragment and was cloned into pGK53 that was linearized with the same enzymes. The new construct was designated pTK-LucR (8931 bp), shown in Figure 2.

15

Example 2

Insertion of Targeting Sequences

A. Generation of vitronectin targeting vector: The targeting construct pTKLR-Vn was generated by inserting vitronectin (VN) DNA sequences into pTK-LucR vector.

Vitronectin (VN) is an abundant glycoprotein present in plasma and the extracellular matrix of most tissues. In a previous study, it was shown that heterozygous mice carrying one normal and one null VN allele and homozygous null mice completely deficient in vitronectin demonstrate normal development, fertility, and survival. This suggests that VN is not essential for cell adhesion and migration during normal mouse development (Zheng, X., et al., Proc Natl Acad Sci U S A 1995 92:12426-30). Mouse vitronectin genomic DNA sequence of 5004 bp was obtained from GenBank database (Accession number X72091). Based on this sequence, a 1.63 kb 3' end vitronectin fragment was amplified (reverse primer, VN1R, SEQ ID NO:11:

25

CTGTATTTAAATCTGCCCCACCCTATTCAGGACAGTAGTC; forward primer, VN1F, SEQ ID NO:12: CCAATGCATCAACCCAGCCAGGAGGAGTGCG) using mouse C57BL/6 genomic DNA as template (Genome Systems, Inc., St. Louis, MO).

30

This fragment was digested with *SwaI* and *NsiI* and cloned into pTK-LucR (linearized with *SwaI* and *SbfI*). This construct was designated as pTK-LucR3. Subsequently, a 2.35 kb 5'end vitronectin fragment was amplified (reverse primer, VN2R, SEQ ID NO:13: AACGCGTCGACTTCGGAGATGTTTCGGGGATAACCAGG; forward primer, VN2F, SEQ ID NO:14: TTGGCGCGCCCCATAGAGAAGAGACACCAAAGGCACGCTC) using mouse C57BL/6 genomic DNA as template. This fragment was digested with *Sall* and *AscI* and cloned into pTK-LucR vector that was linearized with *Sall* and *AscI*. This construct was designated as pTKLR-Vn. Figure 3 shows the restriction map of pTKLR-Vn vector. The polylinker between the neomycin gene and red luciferase gene is used to insert the VEGF promoter or other promoters of interests. The predicted homologous recombination between pTKLR-Vn and vitronectin gene is illustrated in Figure 3A. Upon insertion of the VEGF-LucR transgene cassette, the endogenous vitronectin gene is destroyed. Figure 3B shows the genomic DNA sequence of VN.

15

B. Generation of Fos targeting vector: The targeting construct pTKLG-Fos was generated by inserting FosB DNA sequences into pTK-LucYG vector.

FosB is one of the members of the Fos family. It plays a functional role in transcriptional regulation. It has been shown that FosB mice are born at a normal frequency, are fertile and present no obvious phenotypic or histologic abnormalities (Gruda et al (1996) *Oncogene* 12:2177-2185). A 28.8 kb genomic region that contains mouse FosB DNA sequence was obtained from GenBank database (Accession number AF093624).

Using this sequence, a 1.71 kb 5'end FosB fragment was amplified (forward primer, FosB1F, SEQ ID NO:15: CTGTATTTAAATCCCGTTTCTCACTGTGCCTGTGTC; reverse primer, FosB1R, SEQ ID NO:16: GTCTCCTGCAGGCTTCCTCCTCCTTGTCCTTGCG) using mouse C57BL/6 genomic DNA as template. This fragment was digested with *SwaI* and *SbfI* and cloned into pTK-LucYG vector that was linearized with *SwaI* and *SbfI*. This construct was designated as pTK-LucYG3. Subsequently, a 1.58 kb 3'end FosB fragment was amplified (forward primer, FosB2F, SEQ ID NO:17: AACGCGTCGACGGATGGGATTGACCCCCAGCCCTC; reverse primer, FosB2R,

SEQ ID NO:18: TTGGCGCGCCCCTTGCCTCCACCTCTCAAATGC) using mouse C57BL/6 genomic DNA as template. This fragment was digested with *Sall* and *AscI* and cloned into pTK-LucYG vector that was linearized with *Sall* and *AscI*. This construct was designated as pTKLG-Fos (Figure 4). The polylinker between the neomycin gene and red luciferase gene is used to insert the VEGFR2 promoter (Example 3, Figure 5A), Tie2 promoter (Example 3, Figure 5B), as well as, other promoters of interests. The predicted homologous recombination between the targeting vector bearing the VEGFR2 promoter (Figure 5A) or the Tie2 promoter (Figure 5B) and FosB gene is also illustrated. As shown in the Figures, the VEGFR2-LucYG transgene cassette and Tie2-LucYG transgene cassette is inserted downstream of FosB gene translational stop signal. Therefore, the targeted transgenic mice should still have a functional FosB gene while expressing the transgenes. Figure 4B shows the DNA sequence of FosB.

Example 3

Insertion of Promoter Sequences of Interest

A. pTKLR-Vn/VEGF: Mouse VEGF genomic DNA sequence of 2240 bp that contains a partial VEGF promoter region was obtained from GenBank (accession number: U41383). Accordingly, primers were designed to amplify a 0.69 kb (VF1-VR1A; Table 1) and a 0.98 kb fragment (VF2-VR2; Table 1). It was confirmed that each pair of primers can amplify the predicted product using mouse129SvJ genomic DNA as template.

Table 1

Name	Seq Id No	Sequence
VF1	19	ACCTC ACTCT CCTGT CTCCC CTGAT TCCCA A
VR1A	20	GCTCT GGC GG TCACC CCCAA AAGCA
VF2	21	CCCTT TCCAA GACCC GTGCC ATTTG AGC
VR2	22	ACTTT GCCCC TGTCC CTCTC TCTGT TCGC
KF1	23	GCTGC GTCCA GATTT GCTCT CAGAT GCG
KR1	24	TTCTC AGGCA CAGAC TCCTT CTCCG TCCCT
KF2	25	CAGAT GGACG AGAAA ACAGT AGAGG CGTTG GC
KR2	26	GAGGA CTCAG GGCAG AAAGA GAGCG
TF3	27	AGCTT AGCCT GCAAG GGTGG TCCTC ATCG
TF2	28	CAAAT GCACC CCAGA GAACA GCTTA GCCTG C
TR1	29	GCTTT CAACA ACTCA CAACT TTGCG ACTTC CCG

Conditions for PCR amplification are shown in Figure 6. These primers were used for PCR screening of mouse 129/SvJ genomic DNA BAC (bacterial artificial chromosome) library (Genome Systems, Inc., St. Louis, MO). The library, on average, contained inserts of 120 kb with sizes ranging between 50 kb to 240 kb. A large genomic DNA fragment that contained VEGF promoter region was obtained. Southern blot analysis was performed to map the VEGF promoter region. A unique HindIII restriction site was mapped approximately 7.8 kb upstream of the ATG translational start codon of the VEGF gene. The sequences between HindIII and ATG translational start codon are inserted into the polylinker of pTKLR-Vn vector to finish the construction of targeting vector that contains VEGF-LucR transgene (Figure 3A).

15 B. pTKLG-Fos/VEGFR2

A mouse VEGFR2 genomic DNA sequence of 1079 bp which contains a partial VEGFR2 promoter region was published previously (Ronicke, et al., (1996) *Cir. Res.* 79:277-285). Accordingly, primers were designed to amplify a 0.45 kb (KF1-KR1A; Table 1) and a 0.58 kb fragment (KF2-KR2; Table 1). It was confirmed that each pair of primers can amplify the predicted product using mouse129SvJ genomic DNA as

template. DNA sequences for these primers are shown in Table 1 above and PCR amplification conditions are shown in Figure 6. These primers were used for PCR screening of mouse 129/SvJ genomic DNA BAC library. From the screening, a BAC clone that contained a large genomic DNA fragment was obtained. Based on the published restriction map of VEGFR2, a 4.5 kb HindIII-XbaI fragment that covers the VEGFR2 promoter region was subcloned from the VEGFR2 BAC clone into the pBluescriptSK vector (Stratagene, La Jolla, CA) that was linearized with HindIII and XbaI. The VEGFR2 promoter sequences of 4.1 kb, spanning from a HindIII site to the ATG translational start codon, are subcloned into the polylinker of pTKLG-Fos vector to construct the targeting vector that contains VEGFR2-LucYG transgene (Figure 5A).

C. pTKLG-Fos/Tie2

Mouse Tie2 genomic DNA sequence of 477 bp which contain a partial Tie2 promoter region has been published previously (Fadel et al (1998) *Biochem. J.* 330(Pt. 1):335-343). Accordingly, primers were designed to amplify a 0.45 kb (TF3-TR1; Table 1) and a 0.47 kb fragment (TF2-TR1; Table 1). It was confirmed that each pair of primers amplified the predicted product using mouse129SvJ genomic DNA as template. DNA sequences for these primers are shown in Table 1 above. PCR amplification conditions are shown in Figure 6. These primers are used for PCR screening of mouse 129SvJ genomic DNA BAC library. From the screening, a BAC clone containing a large genomic DNA fragment of the Tie2 promoter region was obtained. Based on the published Tie2 genomic DNA restriction map (see, Dumont et al., *supra*), a 10.5 kb Asp718-EcoRV fragment containing the Tie2 promoter region was subcloned ifrom the Tie2 BAC clone into the pSK vector that was linearized with Asp718 and EcoRV. The Tie2 promoter sequences of about 6.8 kb, spanning from the Asp718 site to the ATG translational start codon is subcloned into the polylinker of pTKLG-Fos vector to construct the Tie2-LucYG targeting vector (Figure 5B).

Example 4**Generation of Transgenic Mice Carrying the Constructs of the Present Invention**

5 **A. General Procedure:** Figure 7 depicts a generalized description of generation of transgenic mice using the targeted transgenic vectors described in Example 3. Details regarding embryonic stem (ES) cell culture, transfection, blastocyst injection and implantation to a pseudopregnant foster are described, for example, in Hogan et al (1994) "Manipulating the Mouse Embryo, A Laboratory Manual. Second Edition", Cold Spring
10 Harbour Laboratory Press.

After construction the targeted transgenic construct are transfected into C57BL/6 embryonic stem (ES) cells. (Genome System Inc., Genome Systems, Inc., St. Louis, MO) through electroporation. The antibiotic G418 is used to select for cells in which the DNA construct containing the Neo gene is integrated, either randomly or by homologous recombination. The nucleoside analog gancyclovir is converted by TK to a cytotoxic
15 derivative. DNA that has integrated by homologous recombination lose the TK gene and are resistant to the drug, whereas cells that have incorporated the DNA randomly are likely to retain the TK gene. Thus, cells containing random integrations into a chromosomal location that allows the expression of the TK gene are killed. The G418
20 and gancyclovir resistant clones are then be screened by PCR and Southern blot analysis and those that have homologous DNA recombination is used for FVB/N blastocyst injection (Genome System, Inc.). Between 4-16 blastocysts are transferred to the uterus of a pseudopregnant foster mother. The pups are typically born 17 days after the transfer. Either random bred mice or F1 hybrid mice make suitable recipients. Females of certain
25 random-bred stocks (*e.g.*, CD1 mice, from Charles River Laboratories) have very large ampullae, which makes oviduct transfer easier. These mice also generally make good mothers. Alternatively, F1 hybrid females (*e.g.*, B6 x CBA F1) can be used as recipients. Although their ampullae are smaller, make exceptionally good mothers, rearing litters as small as two pups. See, for example, Hogan et al. (1994), *supra*.

B. Screening for homologous DNA recombination positive ES cells

- 1) **pTKLG-Fos/VEGFR2:** Analysis of homologous DNA recombination between pTKLG-Fos/VEGFR2 targeting vector and the FosB gene is carried out using Southern blot analysis as shown in Figure 8. Genomic DNA prepared from G418 resistant ES cells is digested with PvuII and probed with probe A to confirm the 5' end DNA recombination. PvuII digestion of DNA bearing homologous recombination reveals two separate bands of 8.2 and 4.0 kb, whereas digestion of DNA from homologous recombination negative clones reveals only the 8.2 kb band. The 3' end of DNA recombination is tested by hybridizing NotI digested DNA with probe B. NotI digestion of DNA bearing homologous recombination will reveal two separate bands of >8.2kb and 5.0 kb, whereas digestion of DNA from homologous recombination negative clones will only reveal the >8.2 kb band. Once homologous DNA recombination is confirmed, positive clones are selected for FVB/N blastocyst injection.
- 2) **pTKLG-Fos/Tie2:** Analysis of homologous DNA recombination between pTKLG-Fos/Tie2 targeting vector and the FosB gene is analyzed by Southern blot in a similar manner as described above for pTKLG-Fos/VEGFR2. Once homologous DNA recombination is confirmed, positive clones are selected for FVB/N blastocyst injection.
- 3) **PTKLR-Vn/VEGF:** Analysis of homologous DNA recombination between pTKLR-Vn/VEGF targeting vector and the vitronectin gene is analyzed by PCR. DNA primers designed according to the predicted homologous recombination, are listed in Table 2.

Table 2
PCR primers for analysis of homologous DNA recombination between pTKLR-Vn/VEGF targeting vector and the vitronectin gene

5

5'end primers	
F51 5'- CCCAGTGTCTCTGATTAGGGAGAGCACCTGAG -3'	SEQ ID NO:30
R51 5'- CCAGACTGCCTTGGGAAAAGCGCCTC -3'	SEQ ID NO:31
F52 5'- CAGTGAGAGTCTTCTCTGTCCCTCAATCGGTTCTG -3'	SEQ ID NO:32
R52 5'- TGGATGTGGAATGTGTGCGAGGCCAG -3'	SEQ ID NO:33
3'end primers	
F31 5'- AATCAAAGAGGCGAACTGTGTGTGAGAGGTCC -3'	SEQ ID NO:34
R31 5'- CGGCTCCCCAAAATGTGGAAGCAAGC -3'	SEQ ID NO:35
F32 5'- GAATCCATCTTGCTCCAACACCCCAACATC -3'	SEQ ID NO:36
R32 5'- CGCCTCCTCTCCCCAGTCTCCCCTTG -3'	SEQ ID NO:37

Primers F51-R51 and F52-R52 amplify a 1799 bp and a 1841 bp DNA fragment respectively from the 5'end of the transgene that is integrated into the vitronectin site through homologous DNA recombination, whereas primers F31-R31 and F32-R32
10 amplify a 3549 bp and a 3428 bp DNA fragment respectively from the 3'end of the transgene that is integrated into the vitronectin site through homologous DNA recombination. Clones that allow successful amplification of both the 5'end and 3'end of the integrated transgene are selected for FVB/N blastocyst injection.

15 C. Analysis of chimeric mice

The pups developed from injected blastocysts contain chimeras, as can be identified by their agouti coat color when an ES cell derived from a mouse having a dark coat color (e.g., C57BL/6) is injected into the blastocyst of a light coat color animal (e.g., FVB/N, genotype B/B). DNA analysis (e.g., Southern blotting, PCR) is conducted to
20 further confirm the presence of the transgene in these pups as described above in Section B. These animals may be obtained commercially, for example from The Jackson Laboratory, Bar Harbor, MN.

D. Generating targeted transgenic C57BL/6 mice with white coat color

Breeding of the chimeric mice generates homozygous targeted transgenic mice, as depicted in Figure 9. The targeted mice are used to monitor gene expression through the measurement of luciferase mediated light emission from the mice. In a preferred embodiment, the targeted mouse has a light coat color (e.g., white coat color), because the black colored coat (an example of a dark coat color) of C57BL/6 mice can absorb light emitted from the body and may interfere the sensitivity of the bioluminescence assay. An inbred mouse strain C57BL/6-Tyr C2j/+ strain (Jackson Laboratory, Bar Harbor, MN) is available for this purpose. This strain of mice have white color coat, yet they still have the same genetic background as C57BL/6 mice except that the gene responsible for the black coat color is mutated. Unfortunately, C57BL/6-Tyr C2j/+ ES cells are not currently available. Therefore, the designed breeding program illustrated in Figure 9 is aimed to generate mice that are homozygous for the target transgene and have white coat color. C57BL/6 ES cells are prepared as described above and introduced into a suitable blastocyst (e.g., from the FVB/N strain of mice). The blastocysts are implanted into a foster mother. Chimeric mice are shown in Figure 9 as white animals with black and green patches. Chimeric animals are bred with C57BL/6-Tyr C2j/+ mice to create F1 hybrids. Subsequent breeding of the F1 hybrids generates several type of mice, including the one that is homozygous for the target transgene and has a white coat color (shown in Figure 9 as b/b; L/L), which is used for *in vivo* gene regulation monitoring.

A C57BL/6 mouse and a C57BL/6-Tyr C2j/+ mouse are considered to be substantially isogenic. Accordingly, the method of the present invention exemplified in Figure 9 provides a means for generating breeding groups of substantially isogenic mice in a selected genetic background carrying at least one transgene of interest.

E. Dual luciferase targeted transgenic mice

As described above, two targeting vectors are generated. PTKLR-Vn carries a red luciferase gene and is targeted into vitronectin locus. PTKLG-Fos carries a yellow-green luciferase gene and is targeted into FosB locus. A number of promoters, including VEGF promoter, VEGFR2 promoter, and Tie2 promoter are cloned into these vectors, as described above. Subsequently three type of targeted transgenic mice are generated.

VEGF mice carry VEGF promoter-red luciferase transgene (VEGF-LucR) integrated into vitronectin locus. VEGFR2 mice carry VEGFR2 promoter-yellow-green luciferase (VEGFR2-LucYG) transgene integrated into FosB locus. Tie2 mice carry Tie2 promoter-yellow-green luciferase (Tie2-LucYG) transgene integrated into FosB locus. Through a breeding program illustrated in Figure 10, dual luciferase targeted transgenic mice are produced, carrying both of the VEGF-LucR and the VEGFR2-LucYG transgenes. The degradation of luciferin by yellow-green luciferase and red luciferase generates lights that emit at 540 nM and 610 nM respectively. These wavelengths of light are measured individually using a photo-counting camera (intensified CCD). Therefore, both VEGF expression and VEGFR2 expression, for example, can then be monitored in the same mouse at the same time.

As is apparent to one of skill in the art, various modification and variations of the above embodiments can be made without departing from the spirit and scope of this invention. These modifications and variations are within the scope of this invention.

CLAIMS

What is claimed is:

1. A vector for use in generating a transgenic non-human mammal, said mammal having at least one single-copy, non-essential gene in its genome, said vector comprising

(a) a first selectable marker and a reporter expression cassette, said reporter expression cassette comprising a transcriptional promoter element operably linked to a light-generating protein coding sequence, and

(b) targeting polynucleotide sequences homologous to a single-copy, non-essential gene in said mammal's genome, said targeting polynucleotide sequences flanking (a), wherein (i) the length of the targeting polynucleotide sequences are sufficient to facilitate homologous recombination between the vector and the single-copy, non-essential gene, and (ii) said transcriptional promoter element is heterologous to the single-copy, non-essential gene.

2. The vector of claim 1, wherein said first selectable marker provides a positive selection.

3. The vector of claim 2, wherein said first selectable marker is selected from the group consisting of neomycin phosphotransferase II, xanthine-guanine phosphoribosyltransferase, hygromycin-B-phosphotransferase, chloramphenicol acetyltransferase, and adeninephosphoribosyl transferase.

4. The vector of claim 3, wherein the first selectable marker is neomycin phosphotransferase II.

5. The vector of any of claims 1-4, further comprising a second selectable marker, wherein at least one target polynucleotide sequence is located between said second selectable marker and (a).

6. The vector of claim 5, wherein said second selectable marker provides a negative selection.

7. The vector of claim 6, wherein said second selectable marker is selected from the group consisting of adenosine deaminase, thymidine kinase, and dihydrofolate reductase.

8. The vector of any of claims 1-7, wherein the transcriptional promoter element is selected from the group consisting of an inducible promoter, a repressible promoter, and a constitutive promoter.

9. The vector of claim 8, wherein the transcriptional promoter element is selected from the group consisting of VEGF, VEGFR, and TIE2.

10. The vector of any of claims 1-9, wherein the sequences encoding the light-generating protein are obtained from either procaryotic or eucaryotic sources.

11. The vector of claim 10, wherein the light generating protein is a luciferase.

12. The vector of any of claims 1-11, wherein said reporter expression cassette further comprises other control elements.

13. The vector of claim 12, wherein said control elements are selected from the group consisting of transcription enhancer elements, transcription termination signals, polyadenylation sequences, sequences for optimization of initiation of translation, and translation termination sequences.

14. The vector of any of claims 1-11, wherein said vector is circular.

15. The vector of claim 14, wherein said vector contains at least one restriction site whose cleavage results in a linear vector having the following arrangement of elements: targeting polynucleotide sequence - (a) - targeting polynucleotide sequences.

16. The vector of claim 5, wherein said vector is circular.

17. The vector of claim 16, wherein said vector contains at least one restriction site whose cleavage results in a linear vector having the following arrangement of elements: target polynucleotide sequence - (a) - targeting polynucleotide sequences - (second selectable marker).

18. The vector of any of claims 1-17, wherein the coding sequences of the reporter expression cassette comprise codons that are optimal for expression in a host system into which the expression cassette is to be introduced.

19. The vector of claim 18, wherein said mammal is a rodent.

20. The vector of claim 19, wherein said mammal is a mouse.

21. The vector of any of claims 1-20, wherein said targeting polynucleotide sequences from single-copy, non-essential genes are selected from the group consisting of vitronectin, *fosB*, and galactin 3.

22. A method of producing a transgenic, non-human mammal, said mammal having at least one single-copy, non-essential gene in its genome, comprising
transfecting an embryonic stem cell of said mammal with a linear vector comprising

(a) a first selectable marker and a reporter expression cassette, said reporter expression cassette comprising a transcriptional promoter element operably linked to a light-generating protein coding sequence, and

(b) targeting polynucleotide sequences homologous to a single-copy, non-essential gene in said mammal's genome, said targeting polynucleotide sequences flanking (a), wherein (i) the length of the polynucleotide sequences are sufficient to facilitate homologous recombination between the vector and the single-copy, non-essential gene, and (ii) said transcriptional promoter element is heterologous to the single-copy, non-essential gene;

selecting embryonic stem cells which each have said first selectable marker and reporter expression cassette integrated into its genome;
injecting said embryonic stem cells into a host embryo,
implanting said embryo in a foster mother,
maintaining said foster mother under conditions which allow production of an offspring that is a transgenic, non-human mammal carrying said reporter expression cassette.

23. The method of claim 22, wherein said offspring is capable of germline transmission of said reporter expression cassette.

24. The method of claim 23 further comprising breeding said offspring with a mammal, wherein the mammal is substantially isogenic with the embryonic stem cells, wherein said breeding yields transgenic F1 offspring carrying said reporter cassette.

25. The method of claim 24, further comprising breeding a first F1 offspring carrying said reporter cassette with a second F1 offspring carrying said reporter cassette, wherein said breeding yields transgenic F2 offspring carrying said reporter cassette.

26. The method of any of claims 22-25, wherein said mammal is a mouse.

27. The method of claim 26, wherein said embryonic stems cells are derived from a mouse having a dark coat color.

28. The method of claim 27, wherein said mammal substantially isogenic with the embryonic stem cells has a light coat color.

29. The method of claim 28, wherein said F2 offspring carrying said reporter cassette has a light coat color.

30. The method of claim 29, wherein said embryonic stems cells are derived from a C57BL/6 mouse having a dark coat color, and said mammal substantially

isogenic with the embryonic stem cells is a C57BL/6-Tyr C2j/+ mouse having a light coat color.

31. The method of any of claims 22-30, wherein the light generating protein is a luciferase.

32. A transgenic, non-human mammal, comprising at least one single-copy, non-essential gene in its genome, wherein (i) at least a portion of at least one single-copy, non-essential gene is replaced by polynucleotide sequences heterologous to the gene, such that the gene cannot produce a functional gene product, and (ii) said polynucleotide sequences comprise a first expression cassette which has been introduced into said mammal or an ancestor of said mammal, at an embryonic stage, said first expression cassette comprising

a first selectable marker,

a first transcriptional promoter element heterologous to the gene, and

light-generating protein coding sequences, wherein said light-generating protein coding sequences are operably linked to said promoter element.

33. The transgenic, non-human mammal of claim 32, wherein the single-copy, non-essential gene is selected from the group consisting of vitronectin, *fosB*, and galactin 3.

34. The transgenic, non-human mammal of any of claims 32-33, wherein the first selectable marker is selected from the group consisting of neomycin phosphotransferase II, xanthine-guanine phosphoribosyltransferase, hygromycin-B-phosphotransferase, chloramphenicol acetyltransferase, and adeninephosphoribosyl transferase.

35. The transgenic, non-human mammal of any of claims 32-34, wherein the first transcriptional promoter element is selected from the group consisting of an inducible promoter, a repressible promoter, and a constitutive promoter.

36. The transgenic, non-human mammal of claim 35, wherein the first transcriptional promoter element is selected from the group consisting of VEGF, VEGFR, and TIE2.

37. The transgenic, non-human mammal of any of claims 32-36, wherein said expression cassette further comprises other control elements.

38. The transgenic, non-human mammal of claim 37, wherein said control elements are selected from the group consisting of transcription enhancer elements, transcription termination signals, polyadenylation sequences, sequences for optimization of initiation of translation, and translation termination sequences.

39. The transgenic, non-human mammal of any of claims 32-38, wherein said non-human mammal comprises a second single-copy, non-essential gene in its genome, wherein (i) at least a portion of the second single-copy, non-essential gene is replaced by polynucleotide sequences heterologous to the second gene, such that the second gene cannot produce a functional gene product, and (ii) said polynucleotide sequences comprise a second expression cassette which has been introduced into said mammal or an ancestor of said mammal, at an embryonic stage, said first expression cassette comprising

a second selectable marker,

a second transcriptional promoter element heterologous to the second gene, and light-generating protein coding sequences, wherein said light-generating protein coding sequences are operably linked to said second promoter element.

40. The transgenic, non-human mammal of claim 39, wherein the first transcriptional promoter element in the first expression cassette is different from the second transcriptional promoter element in the second expression cassette.

41. The transgenic, non-human mammal of claim 40, wherein the light-generating protein in the first expression cassette can produce a different color of light relative to the light-generating protein in the second expression cassette.

42. The transgenic, non-human mammal of any of claims 32-41, said mammal being a rodent.

43. The transgenic, non-human mammal of 42, said rodent being a mouse.

44. The transgenic, non-human mammal of any of claims 32-43, wherein the sequences encoding the light-generating protein are obtained from either procaryotic or eucaryotic sources.

45. The transgenic, non-human mammal of claim 44 wherein the light-generating protein is a luciferase.




FIG. 1




FIG. 2




FIG. 3A




FIG. 3B

1 TCACCCACC TGTTCCTCAG GTCCCCCGCC TTCTCAGTGA ACTTCATGTT TAAGAGAGCC TCACCCCGCC AGGTTGTGTT GCCACAGAG GAGGGTGTCT
AGTGGGTTGG ACAAGAGAGT CAGGGCCCGC AGGATCAAT TGAAGTACCA ATTTCCTGGG AGTGGGCGCC TCACACACCA GGTGTCTTC CTGCCACGA

101 CCCACAGCC CCCAGTGTCT CTGATTAGG GAGAGCACT GAGCCCACTG AGAGTCTTCT CTGTCCCTCA ATCCGTTCTG AATTCGCCA CTTCGCCCTC
GGTGTGTGG GGTTCACAGA GACTAAATCC CTCCTGTGGA CTCGGGTGAC TCTCAGAGGA GACAGGGAT TAGCCAGAGC TTAAAGGGT GAACCGGAGG

201 TTATCCAGG GACAGGGCTG CCCACCTAT TCAGGACAGT AGTCTTAAC TGTAGGCCAA CAGACTTTTT ATTGGGCTGG GAGAGAGAGA TGAGGGTCTT
AATAGGTGCC CTGTCCCGAC GGTGGGATA AGTCTGTCA TCAGAAATTG AGCATCGGTT GTCTGAJJAA TAACCCGACC CTCCTTCTCT ACTTCAGAGA

301 GAGCTCAGC CGAGTGGCT CTGATTCCTA CTCTCAGAG GTCCCGCAGC CCAGCCATA CTGAGCAATG GAGGTTGGTT AGGAGAGTT CACAGAGTCC
CTTCGAGTGG GCTCAGCCGA GACTAAGGAT GAGAGTCTCT CAGCCCGTCTG GGTGGGTAT GACTGTGTAC CTCCACCCCA TCCTCTCA

401 ACTCCCGGG TTCTAAGGTT GACTGGGTAG TATTGTCTG AAGAGAAAG TGAJJAAAG GTTATGTGAG ATTCTGCCG ATCTCTTCA CTGGTCCAA
TGAGCGGCC AAGATCCAA CTGAGCCATC ATAAACAGAC TTCTTTCTT ACCTTTTTC CAATACACTC TAGAGCGAG ATCTCTTCA CTGGTCCAA
GACCGGGT

501 GAGGATAAA GCTTTTCTCT CAGAGGGAA AGTAACTMC CACGAGCAG ATATGTCTAC CATCTACAGC CTGTGTTCAG CACCCAGGA CCAAGAGCTG
CTTCTATTT CCGAAJAGA GTCTTCCCTT TACTTGTAG GTGGTTCGT ATATACAGT GATAGTGTCC GACACAGTCT GTGGGTCCCT GGTCTGTGAC

601 CAGCCAGCC CTAGCCAAA CCAGTCTAG GAGTAGAAG GGGTCCAC CTCCAGAGAA GAATAGAGC CTCTGAATGG GCTGGAGCT GGCAGGTACA
GTCCGTTCG GTCAGATTC CTCACTTCTT GTCAGTCTCT CCGAGGTCTG GGTGGGTAT GACTGTGTAC CTCCACCCCA TCCTCTCA

701 AGCCAGTCA TATCATATC ATAGTTTGT TAGTTTCTA GGGCTCTC CTCCCTGGAG AACAAJAGA ACCAGATTGA AGGTGATGA CAGCGGAGT
TGGTCAGT ATAGATATG TATCAACAC ATCCAGGAT CCGGTGAGAG GAGCGACTC TTGTCTCTCT TGTCTACT TGTCTACTT GCTGCCCTA

801 TGAGCTCTG GCTGGGCTG TGGCCAGCC CTGGGGTGA AGTAGAGCC TTGGGCTTC TAGGCTTAGA CTCTCTTTT TTGGCTTGG CAGATGGGA
AGCTCGAGC GAGCCAGAC AGGGTGGG GAGCCGACT TGTATCCG AJAGCCGAG ATCCGATCT CTCTCTTTT TTGGCTTGG GAGATGGGA
GTCTCGAGCT

901 TAGAGAGCA GTAGCTAGA TGGCCCGCC CAGAGCAGC TCCACTTCC CTGGCAGCC ATCCGATTC CCGCTGATG ATTGGGTTT TCTGGCTCA
ATTCTCGT CTACCTATCT AGGGCGGG GTATGTGGC AGGTGAJAG CTGGCAGCC TAGGGTCTAG GGGGCTACT GAGACTTACC GAGCGGAGT

1001 TCTGTACAG CGAAGGGTT AATGCACTG GCAGATCTG GCTTTGATTT CTCCAGCAG GTTGTCTCT TACTATTTA TCTATCTTA TCTATGATC
AGCATTTCT CTTCCCAA TTAGGTGAC GTCTTAGAC CCAJACTTA GAGTCTCT CACAGAGAG ATAGATAAT AGATAGAAAT AGATACATG

1101 TACTATATA TCTATATC TATCTATC TATCTAAT AACTACTAC GTATCTATGT ATCTATCTAT CTATCATCTA CTTACTACT TACTATCTA
ATAGATAT ATAGATAG ATAGATAG ATAGATAG TGTATGATG TGTATGATG ATAGATAG ATAGATAG ATAGATAG ATAGATAG ATAGATAG

1201 CCTATTATT TGTGTGTG AAJAGAACT TTTCTTTGA AACAGATCT TAGCAGTAC CTATGGCTG TTTCGACTC ACTATGAAG CATACTGCG CTCTTACTC
GGATAAJTA ACJAGAACT TTTCTTTGA AACAGATCT TAGCAGTAC CTATGGCTG TTTCGACTC ACTATGAAG CATACTGCG CTCTTACTC
GAGATATG

1301 ACAAGATCC ACTTCCCTG GTCTCTGAT GCTGGGATG AAGCATGT CCAGTACCC CAGCTCCAGT AGGAGCTTGA GAGCAGATTT GCTATGCCTT
TGTTCCTAG TGAGCGACA CAGAGACTCA GAGCTTAAT TTGTGTAC GGTGATGTG GTCCAGGCTA TCTCGAAT CTGTGTAA CAGATCGGA

1401 GCTTAGACA CACACTCAG TCCCGAGCC CCAGCTTCC TGTCTAGAG TTTTCCCAT CTTCTCTCA CTGTATCCCT TGAATCTCT GGGCTCCGA
CGATTTCTT GTTTGAGTC TCCCGAGCC CCAGCTTCC TGTCTAGAG TTTTCCCAT CTTCTCTCA CTGTATCCCT TGAATCTCT GGGCTCCGA
GGGTAGGCT

1501 AJCCCTCAG CCGCCAGCC CTCTCTCTG TGTGTAGGC AAAGTCCAG GTATGGGAT CAAATAGAGC CAGGCTCAT CCCCAGAG GTCCAGAGT
TTGGGAGTC GGGGTGGG GAGGAGAGC ACJAAATCCG TTCTAGGTTT GTATGGGAT CAAATAGAGC CAGGCTCAT CCCCAGAG GTCCAGAGT
GGGTAGGCT

1601 CAAAGTCTG CAGAGCAGAA CAGCTCTGA TGTATGGTT CAGATTTCA GGGCTTCC CTGGAGGCC CCACTATCAG AGCCAGTTT CAGAGAGAG
GTTCTAGAT GTCTCTGTT GTCTAGAGT AGTCTCTCA GTCTAGAGT GGGGCGGG GAGCTTCCG GTGTATGAGT TCGGTCAAA GGTCTCTTC

1701 AGCCAGGCT TGTCTCTCT CCACTAGCA GATCTTCCC CAGAGAGGA GTTCCAAAT GTTCTCCAG CTGTCCGCT GAGCAGAGC AAJGTCTCA
TTCCGTGGA ACAGAGAGG GTCTCTCTC CCACTAGCA GATCTTCCC CAGAGAGGA GTTCCAAAT GTTCTCCAG CTGTCCGCT GAGCAGAGC AAJGTCTCA
TTCCGTGGA

1801 AACAGGCTG ACAGAGAGT GCTTGGGCT TCTCTCTG TGGTGTGCT CTGAJATG TACTCCAGT ACTGCTTCC TGAAGAGAG ACTCTCTCT TGTGTAGAG
TTGTCCGAC TGTCTCTGA CCGAGAGGT AGGAGAGAG TCCAGAGAG GACTTGAAG ATAGAGGTCA TGAAGAGAG ACTCTCTCT TGTGTAGAG

1901 ATCAGAGAG ATCTAGCAA GCGAGAGAG AATCATGAA TAGACAGAG ACTCCAGC CTGCGGCTT CTCTCCAGC CTGATAGCC TTGAGAGT
TAGTCTCTC TAGACTGTT CCGTCTCTC TAGTACTTT TAGACAGAG ATCTCTCTC TAGAGTGTG GAGCGGGGA GAGGAGGTG GACTCATAGG AACTTCTCA

2001 AGAGCTTTC CCGGCTACT TAAAGTGGG CAGAGAGGC GAGGCTGCA TCAACATTGT CTGTATGCC ACTGAAGCT TCGAGATGT TTGGGAGTA
TCTGGAAAG GGGGCTGAC ATTGCCAGG GTCTCTGCG GAGGCTGCA TCAACATTGT CTGTATGCC ACTGAAGCT TCGAGATGT TTGGGAGTA
TCTGGAGTA

2101 ACCAGGCTC AGGAGCCAT CTTCAJAGG CCACTACTG CTACCTGAA AGACAGAGT CAGAGAGGT AGGACATACC GCTGGCCACA GAGAGAGTCC
TGTGCCAGG TCTGGGTA GAGTTTCC GGTCTACTT CTACCTGAA AGACAGAGT CAGAGAGGT AGGACATACC GCTGGCCACA GAGAGAGTCC
CTGTCTAGG

2201 TATATCTTA ACTGCTCTC ACTCTCTCT GAGTCTCTG ACTCTCTCT CTTCAGAGT CCGCAGCAG TCCATGGCAG CTTTACTCT GCTTACTCT
ATATAGGAT TGAAGAGAG TGAAGAGAG CCGTCTCTG TGAAGAGAG TGAAGAGAG TGAAGAGAG TGAAGAGAG TGAAGAGAG TGAAGAGAG

2301 TAGGCTCTG ACTTGAACA AGTAGGCTT CCGTCTCTG TGAAGAGAG TGAAGAGAG TGAAGAGAG TGAAGAGAG TGAAGAGAG TGAAGAGAG
ATCCAGACA TGAAGATTT TCACTAGAA GGGGCTCTC TGAAGAGAG TGAAGAGAG TGAAGAGAG TGAAGAGAG TGAAGAGAG TGAAGAGAG

2401 TTGGGCTAG CAGGCTCTC TCCCTCTCA TCTACTCAT AGCAGTACT CCGTAGACA GGGGAACTG TGTGAGAGC AGATGAGCT AAGGAGATC
AJCCCATAG GTCCGAGTG AGGAGAGAT AGATAGAGT TGTCTACTC GGGATCTTGT GGGATCTTGT GGGATCTTGT GGGATCTTGT GGGATCTTGT

2501 CACCGGCTC CAGAGCTCT CAGTAGATCA CCGTCTCTG CAAAGAGGA CCGATCTCT AGATCCGTA AGGCTCAAA GGGCTTCCA CTGACAGTT
GCTGGCGTG GTCTGAGAG GTCTCTCAGT GAGGCTTCC CCGATCTCT AGATCCGTA AGGCTCAAA GGGCTTCCA CTGACAGTT
GAGGCTTCC

2601 CTCTCTCTG AACTCAGG GTCCCTTGT CAGTGTGTG GGGCTTCTG ATCTCTCTG GTTCTCTAC TTGAGGCTT GGGGCTCTG GCTGTCTCT
GAGGAGAGC TTGTAGTCC CAGGAGACTA GTCTCTCTG GGGCTTCTG ATCTCTCTG GTTCTCTAC TTGAGGCTT GGGGCTCTG GCTGTCTCT
GAGGAGAGC

2701 AGGATCTAG AGGCTCTG GCTTGAAGT GGGCTCTG CAGAGATTA GTTCAAGG TGAAGAGGT TTCTGGGTT GCACAGGCT GTTGTATTT
TCTAGATCC TTCCAGAGC CCAATCTCA GCGAGGAG GCTCTTAAT CCAATCTCA TGAAGAGGT TTCTGGGTT GCACAGGCT GTTGTATTT
CAACATAC

2801 TTCTCTCTG CCGTCTCTG GTCTCTCTG CCAATCTCA TGAAGAGGT TTCTGGGTT GCACAGGCT GTTGTATTT CAACATAC
AAGAGGGA GTAGTGTAT GTCTCTCTG CCAATCTCA TGAAGAGGT TTCTGGGTT GCACAGGCT GTTGTATTT CAACATAC
GTCTCTCTG

2901 GTCTCTCTG GAGGAGAG CCGAGGAG CCGAGGAG TCTGAGACT CTTGGGCT TGTCTCTCT CAGTGTAGT GCACAGAGC TGTGTAGTA
CAGTGTAGT CCGTCTCTG GAGGAGAG CCGAGGAG TCTGAGACT CTTGGGCT TGTCTCTCT CAGTGTAGT GCACAGAGC TGTGTAGTA
AGATATCAT

FIG. 3C-1

3001 AGTGCAGAGC TGTTCACACT GACACTTCTT GCTGCCCATG AAACCCCTGAG TGCAGCGGCC CTTGCATGAC TCTATGGGAG GGAATATCAG GTTTCAGGCC
TCACTTTTGG AGCAGTGTGA CTGTGAAGAA CGACCGGTAC TTTCGGACTC AGCTTCCUUG GAACGTACTG AGATACCCCTC CCTATATAGTC CAATATGTGGG

3101 CAATCTAGGG CACCTGCCCA ACCTGCACCT CCGTGGGTAC CCACCAATCC CCTCCACAC CTTGCTCAOC CAGAGAAACC CATGCCACCA GCGCTAGTAT
GTTAGATCCC GTTGACGGGT TGGACGTGAA GGGATCCATG GGTGGTTAGG GGAGGGTGTG GAACCACTCG GTCTCTTTGG GTACCGTGGT CCGGATCATA

3201 GAAAAAGGCC CTCAGGGGTG CCATGCCAGG CCTCTACCCC AGGGCTTTGG CAAGCTGGGC GCGAGCTTC TGGATCTCG CTGTCTCTGC TGA AAAAAGA
CTTTTTCGGG GAGTCCCCAC GTTACCGTCC GGAGATCGGG TCCCGGAACC GTTGGACCGG CGCTCGAGG ACCTTAGAGC GACAGGAGCG ACTTTTTTCT

3301 AGCAGACTGA AGAAGAGTTC CTAGTTCCCT GGGTTTCTGC CCTTTATTTG CTCATCTCTT GCGCCAGGCC CATTCGCCCT CTCGAAACAC AGCTGCACCA
TCTCTCTGACT TCTTCTCAGG GATCAGGGGA CCGAAGAGCG GGAATATAAC GAGTAGGAGA CCGGTTCGGG GTACCGGGAG GAGGTTTGTG TCGACGTCTGT

3401 AAGGCTCACA TTCCACAGAC CCCAGCCCCA GGAGAGCTGG GAACAGAGAA ACCCTGCCCA AGACCAAAAT CAGTAGGCTC ACCGGCAGGA GGGATACAC
TTCCCAATGT AAGGCTCTTG GGTTCGGGT CTCTCGAC CCATTTCTTT TGGAGCGGT TCTGTTTCA GTCATCCAG TCGCGTCTCT CCTATTTGTG

3501 GCTTAGCTTA GCTGGGGAG TGGAAAGAG CATGTGTGT CACCTCTGA GCGAGTCCCG TTAATCTCC TTAGCTTAC TTTTNTAA GTGGACCAT
CGAATCGAAT CGACCCCTCC ACCTTCTTC GTACCAACA GTGGAGACT GGTTCAGGCC AATTAGAGG ACTCGAATG TTTTNTAA AACCTGTAT CACCTGTGTA

3601 GGTGCTTTC CACCGAAGG GTTAGAGAT TCCGTAGCT AGACAGAGA AAAGGTTTC TCGCTGAGT AGCTTCCAC TCATTCCTAT AAGCGTAT
CCACCGAAGG GAGTAGTCCA CAATCTCTA AGGCACTGA TTTCATCTT TGTGCAAGC AAGGACTCA TGGAGGTTG AGTAAGGTA TCGGCAATA

3701 CGATTACTG TTTGATCAG CTAGGTGCTT GTCCATCTT ACCCCCCCT TCGAATCTGG ATTTTGGGG CAGAAAGGG GGTGGGGGA GAGCTGCCAA
GCTAAATGAC AAATGATGCC GATCCAGAA CAGGCTAGGA TGGGGGGGA AGCTTAGACC TAAAGACCC GTTCTTCCCC CCACCCCTCT CTGACCGT

3801 GCACTTTGG GAGGTTTTC TTTTCTCTC ABAAGAGAC AAAGCTTCAT TCTGGGCTC TCTTGTCT CTCTAGCTG GGTGTACAG CAGTAAGAT
GTTGAAAGCC CCGCAAGAG TATTTCTTG ABAAGAGAT TTTTCTCTG TTTGAGAGG TCGAGCGAG AGGAACAGA GAGATTGAC CCACATGTC GTATCTCTCA

3901 AGTGGTCAAG AGCTATCTT TCTTCTTCA TTTTCTTCA TTTTATTTT TTAATTTT GTGTAAAT GTCTCTCAG AGTGTCTAT GTGCACCA
TCAACCAATC TCAGATAGA AGAAGAAAT AAAAAAATC TAAATAAAT AATACAAA CACATTTCA GTCTCTCAG CAGAGAGTGT GTGCACCA
CAGAGAGTGT TACAGTAGA CAGTGTGT

4001 TGCATGCTT GTGTCTATG AGTTCAGAG AGGCTTTGA ATACCTGGA ACTGAGTTT TGAACATTA TGAGCTGCG TGTGATGCT GAGATCAAA
AGCTACAGAA CAGGATACC TCCAGTCTTC TCCGAAACT TATGGAGCT TGACCTCAA ACTTGTCAAT ACTGAGGCC ACACCTAGCA CTCTAGTTT

4101 CCGAGTCTT CTGTAGAAC AATGACTCTT AAGGCTGAG CCATCTTTCC AGTCCCAAG CCATTCCTG AGGCTTTAC TAAATCAAT ATCTCGGG
GGTTCAGGA GACTTCTTG TTCTAGAA TTTCGACTC GTTAGAAGG GGTAGAGGAC TCCGAAAGT ATTAGGTAC TTAGAGCCC

4201 GACCACTCT GGCACACTT CAATGACCTC ATTTATTTA AAAAAAAT GGAATCAT GGCATCTTT CTAGACTCAG ATACTAAGT GATTCTCT
CTGTGGGAC CCGTGTGAA GTTACTGAG TAAATAAAT TTTTCTTTA CCGTAGTAC CCGTNTGAA GATCTGAGT TATGATCAC CCTAAGAGA

4301 ATAAAGAGT GCTCACTGG GTAGAGTCC AGGTTTGGG CCAATTCOA AGCACTGCA CACTTCTGAA GCGGCTCGT TTTCTGTCT GTATACAG
TATTTCTCA CGAGTGAACC CATCTCAGG TCCAAAGCC GTTTTAGGT TGTGACCGT GTGAAGACTT CCGGGAGCA AAGACAGA CATAGTCT

4401 GCGAGCGTC CTTGTGTGTC TCTTCTCTAT GGCAGCGAGT AGTCTCAGCG CCMAATGAA ACATTAAT TTAATCCCTA CAGACCGTG AAGCTAAGT
CGTCCACAG GAAACACAG AGAGAGATA CCGGCGTCA TCGAGTGGC COTTTTACTT TGTGATTAA AATGAGGAT GTCTGCCAC TTGGATTCA

4501 GGAACCGGC ATTAAGGCG TTGAGAAAT TCACTGCGA TTCTTTAACC ATCGGAGGG GAGGTGGATA CATGTAGCCA GCTTCTTCC ACATTTTGG
CCTTGGCGG TAAATTCAG AATTCCTAG AGTTGACCT AAGAAATGG TAGGCTGCC CTGCACCTAT GTACATCGT CGACAGAGG TGTAAAGCC


4601 GAGCGAGCG AGCGGTAGA AATGAGAGC AGCTCTTAC AGCGCTTCT ACAGCATCTT GCACACACC AAGGGGAGC TGGGAGAGG AGCGGAGCC
CTCGCTGCG TCCCATCTT TTAGCTTCTG TCGAGAAAT TCGGAAAGA TGTGTAGAA CGTGTGTG TTCCCTCTG ACCCTCTCC TCGGCTGGG

4701 AAGTGTGGC GTGGCTGAG AGCTGGGTA GCTTGGGCG TGGTTCGGG GCGAGCGCG TGAACCTAG AGCGGGGGG TCAATCTCT GACTCTGCT
TCCACAGCG CAGCGACTC TCGACCCAT CCGAAGCGG ACCAGCGCC GCGCTGGG ACTTTGATC TCGGGGGG AGTTTAGGA CTGAGAGAC

4801 CTCAGAGCG TGTGTCTGT TTAGCACTT AGCTGGGCT TCGTGAAT GAGCAGCGC TTTGTTCGG GCACCGGCT CTCACCTCT CCGGCTTGG
GAGTCTCGC ACCAGAGCA ACTGTAGAA TCGAGCGAC AGGAATCTA CCTGTGCG AAACAGGCC CTTGGCGCA GAGTGGAG GCGGAGAGC

4901 TCCAGCTTC TCTCTCTT CATGCCCTC CTAGTCTCT GAGTCCCGA GCTGCCCTC TCTTCTCT TCTACACTG TAGGCCAGCA CCTTACCGG
AGTACGAG AGAGAGGAA GTACGGGAG GATTCAGAA CTCAGGCGT CGACCGGAG AGGAGAGCA AGATGTGAC ATCGGTTGT GGAATGCGC

FIG. 3C-2

**FIG. 4A**

1 GCAGCTGGCC AAACCTTGGC GATGCGGCTG CAAAGTATAT ACCGCTGGCT TAGCAGAGCC TGAGAACATT TAGCGGAAAG CGGCTGCTCT AAGCGGAAAG
CGTGGAGCGG TTTCGACAGG CTAGCGGCAC GTTTCATATA TGGGCGACCA ATGCTCTTGG ACTCTTGAAA ATCGCTTTTC GCGCGAGGGA TTGCGCTTCC

101 TAGCGAAGTA GGGGAGAGAA AAGAAJAJAA AATATCCAGA GAAGCTTCCA GAAGCTGCTC CTCTTCCCTC TTCTTTCAAA AGGACTGCAA GTCCCGAAGTC
ATCGGTTCAT CCGCTTCTTT TTCTTATGTT TTCTAGGTTT CTTCGAGAGG GAGAGGGAG AGGAAAGTTT TCTTGAGTTT TCTTGAGTTT CAGCGGTCTAG

201 ACCCTCCACC CAGCAGAGAT TAGGCGCTCG AACCCTGGTC ACCTGCGCTC CGGCTGCTGC CGAACGTAAAC GGGGACCGCG TCGTAAAGCC GTGAGCGGCT
TGGGAGGTGG GTGCTTCTCA ATCGCGGAGC TTGGGGCCAG TCGGACCGAG GCGGAGGAGC GCTTCGATTG CCGCTTGGCC AGCGATTTCG CACTGCGCGA

301 GGAATCTTCC GTCTGACCGC GGGCAGCCAC AGGCGCAGCC CCGTCCGGCC GCGCGCGCGC TGAGGTCCGG GCAGGTTCTA TTTTGAAAG CCGAGGCGAC
CCTTAGGAGG CAGACTCGCC CCGCTGCTGG TCGGCGCTCG CCGTCCGGCC GCGCGCGCGC ACTGCGAGCC GTTCGAGAT TCTGAGTTGC GCGTCCGGTG

401 GTTCTAAGG GAGCGCGCAG CCGTGGCTTG TGATTTGGCT TCGGCGCGAG CTTFAGCCAA TCAGCGTTCC CTTCCTATTT GTGAGGCGTA GCTCCCTTCC
CAACGATTTC CTGCGCGCTC GCACCGAAGC ACTAAGCGAC AGCGCGCGCT GAATTCGGTT AGTTCGAGAG GAGGATGAAA CATCTCGCAT CGAGGGAGGG

501 TTGCTTTTGG TGGTCTTTCG CCGTGGCTTG GTCTCCAGGA GAGAGCTAG GATCTTGTCT GCGATCGGGA CTGTTGTICA CCGCATGCTC TGCGAGGACT
AAGGAAJAJA ACCAAGAGAG GCACGACCGC CAGAGGTTCT CCTCTGATCT CTAGAGACAG CCGTAGCCCT GAGCAACAT GGGGTACAG ACCCTCTCA

601 TGTGTGAGCC TGGTCTTTCG TCAATAGCTA GAGGCTTTTG GCTGAGTGT ACCGCTCTCA AGCGGGAAGT GAGGCGCTCA TCGTCTCAG GCACACATAT
ACACAGCTGG TCGGCTTTCG ACTATTCGAT CTCCGAAAGC GCGACTACAA TCGCGAGAT TCGCGGAGT CTTCGGAGT CTTGAGGCTC GATGTGTATA

701 AGTGTCTCT GAGCTCTAGA CACTGATGCC TTGCGAGGT TTCAACACT AGATGAGCTA GCTTACCGAG AGGCGCGAG GTGCTCTCA AAGGCTCTCC
TGCAGAGGA CTGAGATCT GTGAGTCAAG AAGGCTCCAC AATTTTGTGA TCTACTCGAT GCGATGCTC TCGTCCGCTC CACCGAGAT TTTCAGAGCG

801 CTTCCTTTCG TTCCAGGCT CTGATTGGCC AGGATTTCAG CCGTTCCTTC CGCAGCGCGC CTAGAGTATG TAGGCTCTCA GATTCGAT TTGCGGAAAG
GAGGCTTTCG AAGGCTTTCG GACTAAGCGG TCGGTAAGTC TCGGTAAGTC GATCTCTATCA ATTCGAGAT ATTCGAGAT TTGCGGAAAG

901 GCGCGCGCGC GCGCGCTGAG GAGGCTTCTT GCGGACCGAG ATCTATGCT ACCGCTTCCG CTGCGAGACA GTCTGAGAGA TTCTGCTGT CACTTTTCTC
CGCGCGCGCG CCGCGCTTTC CTGCGAGAGA CCGCTTCTTC TCGGTAAGTC TCGGTAAGTC GAGGCTTCTT CACTTTTCTC AAGGCTTTCG

1001 TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG
ACGGAATGTC AAGTATCTTT GCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG

1101 ACTGCGATCT CCGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG
TGAGCTTTCG CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG

1201 TGTGTGAGCC TGGTCTTTCG TCAATAGCTA GAGGCTTTTG GCTGAGTGT ACCGCTCTCA AGCGGGAAGT GAGGCGCTCA TCGTCTCAG GCACACATAT
ACACAGCTGG TCGGCTTTCG ACTATTCGAT CTCCGAAAGC GCGACTACAA TCGCGAGAT TCGCGGAGT CTTCGGAGT CTTGAGGCTC GATGTGTATA

1301 CCGCGCGCGC ACACCTTTCG GTGAGTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG
GAGGCTTTCG CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG

1401 GTTGAACAGA ATCTCTCTT AAGGCTTTCG TCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG
CAACTTCTCT TAGAGATGTA TTGCTTTCG GAGGCTTTCG TCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG

1501 TGTGCTTCT ACAGGCTTTCG GTGAGTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG
ACAGGCTTTCG GTGAGTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTTCG

1601 GTGAGGCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG
CATCTTTCG CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG

1701 CTATCTCTT CCAATGCGCA GTGAGTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG
GAGTGAAGAA GTGAGTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTTCG

1801 GCTGAGTCT CCAATGCGCA GTGAGTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG
CGAGTGAAG GTGAGTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTTCG

1901 GCTGAGTCT CCAATGCGCA GTGAGTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG
CGAGTGAAG GTGAGTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTTCG

2001 TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG
ACGGAATGTC AAGTATCTTT GCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG

2101 GAGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG
CTGAGTCTT CCAATGCGCA GTGAGTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG

2201 AAGTGAAGT AATTTTTCG CCAATGCGCA GTGAGTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTTCG
TCTACTTTCG TCAATGCGCA GTGAGTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG

2301 TTTGCGGCTT AAGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG
AAGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG

2401 AAGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG
TTTCTCTTTC CCAATGCGCA GTGAGTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG

2501 CCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG
GAGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG

2601 CAGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG
TCAATGCGCA GTGAGTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTTCG

2701 ATGTCAGCT CCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG
TCAATGCGCA GTGAGTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTCTT CCGGCTTTCG GAGGCTTTCG

2801 CTGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG
GAGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG

2901 CCAATCTCT GAGTGGCTCT GCGTGGAAAC TATTTTGTCT TAGTCAATT CCGTGTCTTC TACTTCACT ATCTACAGT CTGCGGAGT TGAGTGGTG
GTGAGTGA CTGAGGAG CCGGCTTTCG ATAAACAGC ATTCAATTA GGAACAGAG CCGTGTCTTC ATGAGTGA TAGATGTCTA GAGGCTTTCG ACTGAGCAG

3001 GCGCGCGCGC AGCGGCTTTC TTTCTCTCT TTTTACCTCA GTGCGAGCGC CCGTGTCTTC CCGTGTCTTC TTTTACCTCA TTTTACCTCA TTTTACCTCA
CGCGGCTTTC TCGGCTTTCG AAGGAGAGAA AATATGAGT GTGCGAGCGC CCGTGTCTTC CCGTGTCTTC TTTTACCTCA TTTTACCTCA TTTTACCTCA

3101 CTGCGGCTTC GAGGCTTTCG GCGGCTTTCG AAGGCTTTCG AAGGCTTTCG AAGGCTTTCG AAGGCTTTCG AAGGCTTTCG AAGGCTTTCG AAGGCTTTCG
GAGGCTTTCG CCGGCTTTCG AAGGCTTTCG AAGGCTTTCG AAGGCTTTCG AAGGCTTTCG AAGGCTTTCG AAGGCTTTCG AAGGCTTTCG AAGGCTTTCG

3201 TTCTCTTCAA AATATATTT CCGTGTCTTC TTTTATATTT CCGTGTCTTC TTTTATATTT CCGTGTCTTC TTTTATATTT CCGTGTCTTC TTTTATATTT
AAGGAGAGTT TTTTATATTT CCGTGTCTTC TTTTATATTT CCGTGTCTTC TTTTATATTT CCGTGTCTTC TTTTATATTT CCGTGTCTTC TTTTATATTT

3301 TCAAGTCTTC TTTTATATTT CCGTGTCTTC TTTTATATTT CCGTGTCTTC TTTTATATTT CCGTGTCTTC TTTTATATTT CCGTGTCTTC TTTTATATTT
AAGGAGAGTT TTTTATATTT CCGTGTCTTC TTTTATATTT CCGTGTCTTC TTTTATATTT CCGTGTCTTC TTTTATATTT CCGTGTCTTC TTTTATATTT

3401 CCGTGTCTTC TTTTATATTT CCGTGTCTTC TTTTATATTT CCGTGTCTTC TTTTATATTT CCGTGTCTTC TTTTATATTT CCGTGTCTTC TTTTATATTT
AAGGAGAGTT TTTTATATTT CCGTGTCTTC TTTTATATTT CCGTGTCTTC TTTTATATTT CCGTGTCTTC TTTTATATTT CCGTGTCTTC TTTTATATTT

3501 CTGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG
GAGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG

FIG. 4B-1

1601 CAGTGGGAG GGAAGTGTCC AGCCGCCCTGG ATCAAGCAACA AGAATGTATG AGTGTGGGTT TGGCGGGTGT AAGCTACTCT GTGTGGTCC TGACCAACAA
GTCAAGCCCTC GCTGACAGAG TGGGGGAGCC TAGTGTGTGT TCCATACATC TCACAGCCCA ACCCGCCAC TTGGATGAGA CACACAGAGG ACTGGTGTGT
1701 TTCTCTTTTC TCTGTCTCTT ATGACCTGGC CCTGCTGGGA TCCTATAGCA AACTGATCAG CTTCAGAGAG AAAAGGCAGA CTTGGAGAGG GAGATTCGCC
AAGAGCAAGG AGACAGAGGA TACTGGACCG GACAGACCTT AAGTAATCTT TTGACTAGTC GAACTTCTCC TTTTCCGTCT CAGCTCAGC CTCTAGCGCC
1801 AGCTOCAMAA AGAGAAAGAA CCGCTGGAGT TTGTCTGTGT GCGCCACAAA CCGGCTGCA AGATCCCTTA CGAAGAGGGG CCGCGCCAG GCGCGCTGCC
TGAGAGTTT TCTCTTCTTT GCGGACCTCA AACAGGACCA CCGGTGTGTT GCGCGGACGT TCTAGGGGAT GCTTCTCCCC GCGCGCGGTC CCGCGGAGCC
1901 CGAGGTGAGA GATTTCGAG GGTCAACATC GGTCAAGGAA GAGGCTTGG CTTGCTCTCT GCGCGCCCTT CCACACCCG CCGTGGCTTT CAGAGCACCC
GCTCCACTCT CTAAGCGGTC CCAATTTGAG GCGATTTCTT GCGATTTCTT TGTGCGAAGC CCAAGGACCA CCGCGGGGGA GTTGTGTGCG GCGACCGGAA GGTCTGTGCG
2001 GAGGAGCCAC CCGCCAGCTT GAGGCTTCTT CTCTTTACAC ACAGTGAAT TCAGTCTCTC TCGGACCCCT TCCGACCCCT TCCGACCCCT TCCGACCCCT
GCTCTGCGTG GCGGTGTGGA CCGCGGAGG CTGCGGAGA GAGAAATGTG TGTCTCTTCA TCAAGTCTCT CCGGACCCCT TCCGACCCCT TCCGACCCCT
2101 GGTGTGTCTT CAGCTGCGCG GAGGTCTGCC GGTGTGCGCG GCGCCACGCC ACCAGCGCCA GCGAGCAGCC GTCCGACCGG CTGAACCTCC CTTGCTCTCT
CACACTTGGG CCACTTCTGG CCGGACCTCC CTGAGAGCC GCGAGTGTGG AGAGCGGCGC CCGCTGTGCG CAGCTGTGCG CAGCTGTGCG CAGCTGTGCG
2201 TGCTCTGTAA ACTCTTAAAG TGAAGAAATCT GTTGTGTTTG TTTGTGTTGG CCGTCTCTCT TCTCTCTCTT TCTCTCTCTT TCTCTCTCTT TCTCTCTCTT
AGAGAGATCT TGAAGAAATCT GTTGTGTTTG TTTGTGTTGG CCGTCTCTCT TCTCTCTCTT TCTCTCTCTT TCTCTCTCTT TCTCTCTCTT TCTCTCTCTT
2301 GTGTGAGAGC TTTGACTCTT CTGTCTGACC AGCTGCGGCC TCTGCGATCG GACATGAGCG AAGGACCTCC TTTGTGTTTG GTCTCTCTCT TCTGTGTTTG
AACTGAGAA TGAAGAAATCT GTTGTGTTTG TTTGTGTTGG CCGTCTCTCT TCTCTCTCTT TCTCTCTCTT TCTCTCTCTT TCTCTCTCTT TCTCTCTCTT
2401 TGTGCGCGCG CGAGACCGGA GAGCTGTGTA CTTTGTGGAC AGCGGTGTGG CCGCGGAGTGA ACACCCCTCC TGTGCGCGCG TGTGCGCGCG TGTGCGCGCG
ACAGCGCGCC GCTCTGCGCT CTGCGACCTT GAAACCCCTG TCCCGGACCC CCGCGGAGTGA ACACCCCTCC TGTGCGCGCG TGTGCGCGCG TGTGCGCGCG
2501 ACTTGTGGGG ATGAGTGGCT GACTGGGTGG GTAGGGTGGG GTCCAGCCCG CAGCTTTGCG GTCTTACGTT AGCTGTGAGG AGCTGTGAGG AGCTGTGAGG
TGAAGACCCG TATCTACCGA CTGAGCCACC CTGAGCCACC CTGAGCCACC CTGAGCCACC CTGAGCCACC CTGAGCCACC CTGAGCCACC CTGAGCCACC
2601 GGTGAGGGGT GGTGTGAGGT GAGCTGCGCA TGCACTTCCA GAGAGACCCA AGAGGAAAT GACAGCAGCC TCTGTCTCTT CTTTTCGCC ACCGACCCAT
CACACTTGGG CCACTTCCA GCTGAGCGGT AGGTGAGGT CTCTCTGAGT CTCTCTGAGT CTCTCTGAGT CTCTCTGAGT CTCTCTGAGT CTCTCTGAGT
2701 CCACTCTCAA GGTGTGAGGT GAGCTGCGCA TGCACTTCCA GAGAGACCCA AGAGGAAAT GACAGCAGCC TCTGTCTCTT CTTTTCGCC ACCGACCCAT
GTGAGGAGTT CCACTTCCA GCTGAGCGGT AGGTGAGGT CTCTCTGAGT CTCTCTGAGT CTCTCTGAGT CTCTCTGAGT CTCTCTGAGT CTCTCTGAGT
2801 GAGCAATGTA GCGCGGAGT GAGGAGGATC GATGCGCGCT TTGGGAGTGT GCTTACCCCA CTCTGCGGTA CTTTTCGCC ACCGACCCAT
CTCTTACCT CTCTTACCT CTCTTACCT CTCTTACCT CTCTTACCT CTCTTACCT CTCTTACCT CTCTTACCT CTCTTACCT CTCTTACCT
2901 TCACTCTTTC CTTGCTGGA AAAGTGTCTT AGGTGAGGT TTTTCTCTCT TCTGCTGAGT GAGAGACCCA AGAGGAAAT GACAGCAGCC TCTGTCTCTT
AGTCAGAAAG GAGGAGGAGT TTTGAGCGAG TCCAGCTTAA TCCAGCTTAA TCCAGCTTAA TCCAGCTTAA TCCAGCTTAA TCCAGCTTAA TCCAGCTTAA
3001 TATTATGCTA TATTATGCTA AAGTGTCTCT AGCTGAGGAG CCACTGAGGAG CCACTGAGGAG CCACTGAGGAG CCACTGAGGAG CCACTGAGGAG
ATTAATGCTA TATTATGCTA AAGTGTCTCT AGCTGAGGAG CCACTGAGGAG CCACTGAGGAG CCACTGAGGAG CCACTGAGGAG CCACTGAGGAG
3101 CCACTTCTCT GAGGAGGAGT ATACTGTGAA TGAATGAGT GATGCTGAGG CCGCGCGGAT GGGATGAGC CCACTTCTCT GAGGAGGAGT
GAGGAGGAG CCACTTCTCT ATACTGTGAA TGAATGAGT GATGCTGAGG CCGCGCGGAT GGGATGAGC CCACTTCTCT GAGGAGGAGT
3201 CCACTTCTCT ACTGCTTCT TGAAGAGAGG AGGAGGAGG ACTGCTCTCT TGAAGAGAGG AGGAGGAGG ACTGCTCTCT TGAAGAGAGG
GAGGAGGAG TGAAGAGAGG AGGAGGAGG ACTGCTCTCT TGAAGAGAGG AGGAGGAGG ACTGCTCTCT TGAAGAGAGG AGGAGGAGG
3301 TTTTCTCTTT AAGTGTCTCT CTTTCTCTCT CTTTCTCTCT CTTTCTCTCT CTTTCTCTCT CTTTCTCTCT CTTTCTCTCT CTTTCTCTCT
AAAGAGAGAA TGAAGAGAGG AGGAGGAGG ACTGCTCTCT TGAAGAGAGG AGGAGGAGG ACTGCTCTCT TGAAGAGAGG AGGAGGAGG
3401 GAGTGTCTTA GCGCTGAGG TTTGCTCTCT CTATCTCTGA TATTGAGATC CCACTTCTCT TTTTGGAGT TTTTGGAGT TTTTGGAGT TTTTGGAGT
CTTCAAAAGT CCGGAGTCTG AAAGGAGGAG GATGAGGAGT ATTAAGATTA GGGTGTATTA AAAGGAGTAT GATGAGGAGT TTTTGGAGT
3501 ATCCACTTCC ATCCACTTCC TTAGGTTAGG TTAGGTTAGG TTAGGTTAGG TTAGGTTAGG TTAGGTTAGG TTAGGTTAGG TTAGGTTAGG
TAGGTTAGG TTAGGTTAGG TTAGGTTAGG TTAGGTTAGG TTAGGTTAGG TTAGGTTAGG TTAGGTTAGG TTAGGTTAGG TTAGGTTAGG
3601 AATATTGGGG AGATGGGCGG TACCGCGCGT CCGCGCGGCT GATGAGGAGT GATGAGGAGT GATGAGGAGT GATGAGGAGT GATGAGGAGT
TGTAAAGGAG TGTAAAGGAG TGTAAAGGAG TGTAAAGGAG TGTAAAGGAG TGTAAAGGAG TGTAAAGGAG TGTAAAGGAG TGTAAAGGAG
3701 CCGCGAGCTA TTTATCTCTT TCTGCTTCT CCAAAAGCAC TTTATCTCTA TTTATCTCTA TTTATCTCTA TTTATCTCTA TTTATCTCTA
GGGCTCTGAT AAATAGGAGG AGGAGGAGG GTTTTCTGTT TTTATCTCTA TTTATCTCTA TTTATCTCTA TTTATCTCTA TTTATCTCTA
3801 GGTGTGGGT GGTGTGGGT GAGCTTCTCT GTTTTCAAGT GTGCTGTGGA GTTCAAAATC CTTTCTGCGG ATTTGAGTCA GACTTCTGCG CTGTCTCTTT
CAGCGACCA CCAAGCCAGG CCAAGCCAGG CCAAGCCAGG CCAAGCCAGG CCAAGCCAGG CCAAGCCAGG CCAAGCCAGG CCAAGCCAGG
3901 TTGTCTCTTT TTTGTGTTTG TCTGCGCTCC TCTGCGCTCT GAGAGCAGTC CCGCGCTCTC CTTTATCTCT TTCTCAAGTC TGTCTGCTCT AGACCACTTC
AACAGTAAA AACAGTAAA TCTGCGCTCC TCTGCGCTCT GAGAGCAGTC CCGCGCTCTC CTTTATCTCT TTCTCAAGTC TGTCTGCTCT AGACCACTTC
4001 CAACTGTCT CCACTCTCAA TGTGAGGAT ACTGAGACTA CTCTGTCTAG CCGGAGGAG GCGGAGGAG GCGGAGGAG GCGGAGGAG GCGGAGGAG
GTGTGAGGAT GTGTGAGGAT ACTGAGACTA CTCTGTCTAG CCGGAGGAG GCGGAGGAG GCGGAGGAG GCGGAGGAG GCGGAGGAG
4101 GGTGTGAGT TTTTCAAT CTATGCTGTT GAGATTTCTG GGTGAGGAT TCTGATCAG AGAGGAGGCT GCGGAGGAGT GCGGAGGAGT GCGGAGGAGT
TGTGAGGAT TTTTCAAT CTATGCTGTT GAGATTTCTG GGTGAGGAT TCTGATCAG AGAGGAGGCT GCGGAGGAGT GCGGAGGAGT GCGGAGGAGT
4201 TAGGCTCTAC ATCGGAGTGT CAGAGCTGTA TTTGTGATTT TTTTCAATTT GTTTTGTGTT ATTTGCGCC TGAAGGAGG GGTGCTGAGT GGTGCTGAGT
ATCGGAGTGT CAGAGCTGTA TTTGTGATTT TTTTCAATTT GTTTTGTGTT ATTTGCGCC TGAAGGAGG GGTGCTGAGT GGTGCTGAGT
4301 CCGCTCTCTCT CTTGCTTCT CCACTGCTCC CAATGAGAG GTTATTTTTC CTTTAAAAA ACTGATCTCT TCAAGTCAAA GTGCTGATTT TCCCTGAGA TCTATCTAT
GGGAGGAGG GGAAGCAGA CTTTCTTCT CTTTCTTCT GGTATGAGG GTTATTTTTC CTTTAAAAA ACTGATCTCT TCAAGTCAAA GTGCTGATTT TCCCTGAGA
4401 GCTTCTCTTT CAGAAAAAG GAGTTTGGAT TCTTGGGTA GTCTTCTCTG CACTTGTGAG GAGGCTTAC GATGCTTCTG GATGCTTCTG GATGCTTCTG
CAGAAAAAG GCTTCTTCT CTTTCTTCT GGTATGAGG GTTATTTTTC CTTTAAAAA ACTGATCTCT TCAAGTCAAA GTGCTGATTT TCCCTGAGA
4501 AGTGAGACT TGTAGGTTT TCCATGTTT CCGGCTGAG CCACTGCTCT GAAAGATTTA GCGGCGGAAA AGTGTAGGT ACCAAATTTT GTGAGGCTC GTGAGGCTC
TCACTCTCTA ACATCTCAA AGGCTCAAAG GGTGCGAGCC GGTGCGAGT GGTGCGAGT GGTGCGAGT GGTGCGAGT GGTGCGAGT
4601 TGGAGAAAT TCAATGCTCG AAAAGATTTT ATTCAGCTTG GGTGTGCAAT GAACTTCTAG CTTGAAATTC CTAAGTCTAG CTAAGTCTAG CTAAGTCTAG
ACCTCTCTAA AGTACTAGCC TTTTCTTAAA TGAAGTGAAC TGAAGTGAAC TGAAGTGAAC TGAAGTGAAC TGAAGTGAAC TGAAGTGAAC
4701 GAAATCTCTA CCAATTTAGA GGTGAGGACA AGGGATCTAA CTGCTGAGT TCAATGCTAT GTGATGCTA GATACCAAGC GCAAGATCTT CCAAGATCTT
CATTEAGAT CCAATTTCTT TACACAGAG GAGCCAGAG TTTGTGTGTT AGGCTGAGT TTTGTGTGTT AGGCTGAGT TTTGTGTGTT AGGCTGAGT TTTGTGTGTT
4801 GAGGCTTGG TACACAGAG CCACTGCTCT TTTGTGTGTT AGGCTGAGT TTTGTGTGTT AGGCTGAGT TTTGTGTGTT AGGCTGAGT TTTGTGTGTT
CTCCGAGCC CTTGAGTCTT CTTGAGTCTT CTTGAGTCTT CTTGAGTCTT CTTGAGTCTT CTTGAGTCTT CTTGAGTCTT CTTGAGTCTT
4901 ATCAGAGT CAGAGCTCT CTTTGAAGC ACTAGAGAGC CCGAGAGT GGTGAGTCTT TTTGAGCAGA GAAGGTTAGA GTTAGGTTCTT CTTCTCTCTT
TACGCTCTCA GTCTCTCAAA AGTCTCTTTC TCAAGTCAAA AGTCTCTTTC TCAAGTCAAA AGTCTCTTTC TCAAGTCAAA AGTCTCTTTC
5001 AGTGAGACT TGTAGGTTT TCCATGTTT CCGGCTGAG CCACTGCTCT GAAAGATTTA GCGGCGGAAA AGTGTAGGT ACCAAATTTT GTGAGGCTC
TCACTCTCTA ACATCTCAA AGGCTCAAAG GGTGCGAGCC GGTGCGAGT GGTGCGAGT GGTGCGAGT GGTGCGAGT GGTGCGAGT
5101 TGGAGAAAT TCAATGCTCG AAAAGATTTT ATTCAGCTTG GGTGTGCAAT GAACTTCTAG CTTGAAATTC CTAAGTCTAG CTAAGTCTAG
ACCTCTCTAA AGTACTAGCC TTTTCTTAAA TGAAGTGAAC TGAAGTGAAC TGAAGTGAAC TGAAGTGAAC TGAAGTGAAC TGAAGTGAAC
5201 GAAATCTCTA CCAATTTAGA GGTGAGGACA AGGGATCTAA CTGCTGAGT TCAATGCTAT GTGATGCTA GATACCAAGC GCAAGATCTT CCAAGATCTT
CATTEAGAT CCAATTTCTT TACACAGAG GAGCCAGAG TTTGTGTGTT AGGCTGAGT TTTGTGTGTT AGGCTGAGT TTTGTGTGTT AGGCTGAGT
5301 GAGGCTTGG TACACAGAG CCACTGCTCT TTTGTGTGTT AGGCTGAGT TTTGTGTGTT AGGCTGAGT TTTGTGTGTT AGGCTGAGT TTTGTGTGTT
CTCCGAGCC CTTGAGTCTT CTTGAGTCTT CTTGAGTCTT CTTGAGTCTT CTTGAGTCTT CTTGAGTCTT CTTGAGTCTT CTTGAGTCTT
5401 ATCAGAGT CAGAGCTCT CTTTGAAGC ACTAGAGAGC CCGAGAGT GGTGAGTCTT TTTGAGCAGA GAAGGTTAGA GTTAGGTTCTT CTTCTCTCTT
TACGCTCTCA GTCTCTCAAA AGTCTCTTTC TCAAGTCAAA AGTCTCTTTC TCAAGTCAAA AGTCTCTTTC TCAAGTCAAA AGTCTCTTTC
5501 AGTGAGACT TGTAGGTTT TCCATGTTT CCGGCTGAG CCACTGCTCT GAAAGATTTA GCGGCGGAAA AGTGTAGGT ACCAAATTTT GTGAGGCTC
TCACTCTCTA ACATCTCAA AGGCTCAAAG GGTGCGAGCC GGTGCGAGT GGTGCGAGT GGTGCGAGT GGTGCGAGT GGTGCGAGT
5601 TGGAGAAAT TCAATGCTCG AAAAGATTTT ATTCAGCTTG GGTGTGCAAT GAACTTCTAG CTTGAAATTC CTAAGTCTAG CTAAGTCTAG
ACCTCTCTAA AGTACTAGCC TTTTCTTAAA TGAAGTGAAC TGAAGTGAAC TGAAGTGAAC TGAAGTGAAC TGAAGTGAAC TGAAGTGAAC
5701 GAAATCTCTA CCAATTTAGA GGTGAGGACA AGGGATCTAA CTGCTGAGT TCAATGCTAT GTGATGCTA GATACCAAGC GCAAGATCTT CCAAGATCTT
CATTEAGAT CCAATTTCTT TACACAGAG GAGCCAGAG TTTGTGTGTT AGGCTGAGT TTTGTGTGTT AGGCTGAGT TTTGTGTGTT AGGCTGAGT
5801 GAGGCTTGG TACACAGAG CCACTGCTCT TTTGTGTGTT AGGCTGAGT TTTGTGTGTT AGGCTGAGT TTTGTGTGTT AGGCTGAGT TTTGTGTGTT
CTCCGAGCC CTTGAGTCTT CTTGAGTCTT CTTGAGTCTT CTTGAGTCTT CTTGAGTCTT CTTGAGTCTT CTTGAGTCTT CTTGAGTCTT
5901 ATCAGAGT CAGAGCTCT CTTTGAAGC ACTAGAGAGC CCGAGAGT GGTGAGTCTT TTTGAGCAGA GAAGGTTAGA GTTAGGTTCTT CTTCTCTCTT
TACGCTCTCA GTCTCTCAAA AGTCTCTTTC TCAAGTCAAA AGTCTCTTTC TCAAGTCAAA AGTCTCTTTC TCAAGTCAAA AGTCTCTTTC
6001 CTAAGAGAGA GACTCTCTCT TTAAGAGAG AATGCTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC
GCTCTCTCTT CTAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC
6101 TTAATGCTAA AATGCTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC
GCTCTCTCTT CTAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC
6201 TAGGCTCTAC ATCGGAGTGT CAGAGCTGTA TTTGTGATTT TTTTCAATTT GTTTTGTGTT ATTTGCGCC TGAAGGAGG GGTGCTGAGT GGTGCTGAGT
ATCGGAGTGT CAGAGCTGTA TTTGTGATTT TTTTCAATTT GTTTTGTGTT ATTTGCGCC TGAAGGAGG GGTGCTGAGT GGTGCTGAGT
6301 CCGCTCTCTCT CTTGCTTCT CCACTGCTCC CAATGAGAG GTTATTTTTC CTTTAAAAA ACTGATCTCT TCAAGTCAAA GTGCTGATTT TCCCTGAGA
GGGAGGAGG GGAAGCAGA CTTTCTTCT CTTTCTTCT GGTATGAGG GTTATTTTTC CTTTAAAAA ACTGATCTCT TCAAGTCAAA GTGCTGATTT
6401 GCTTCTCTTT CAGAAAAAG GAGTTTGGAT TCTTGGGTA GTCTTCTCTG CACTTGTGAG GAGGCTTAC GATGCTTCTG GATGCTTCTG GATGCTTCTG
CAGAAAAAG GCTTCTTCT CTTTCTTCT GGTATGAGG GTTATTTTTC CTTTAAAAA ACTGATCTCT TCAAGTCAAA GTGCTGATTT TCCCTGAGA
6501 AGTGAGACT TGTAGGTTT TCCATGTTT CCGGCTGAG CCACTGCTCT GAAAGATTTA GCGGCGGAAA AGTGTAGGT ACCAAATTTT GTGAGGCTC
TCACTCTCTA ACATCTCAA AGGCTCAAAG GGTGCGAGCC GGTGCGAGT GGTGCGAGT GGTGCGAGT GGTGCGAGT GGTGCGAGT
6601 TGGAGAAAT TCAATGCTCG AAAAGATTTT ATTCAGCTTG GGTGTGCAAT GAACTTCTAG CTTGAAATTC CTAAGTCTAG CTAAGTCTAG
ACCTCTCTAA AGTACTAGCC TTTTCTTAAA TGAAGTGAAC TGAAGTGAAC TGAAGTGAAC TGAAGTGAAC TGAAGTGAAC TGAAGTGAAC
6701 GAAATCTCTA CCAATTTAGA GGTGAGGACA AGGGATCTAA CTGCTGAGT TCAATGCTAT GTGATGCTA GATACCAAGC GCAAGATCTT CCAAGATCTT
CATTEAGAT CCAATTTCTT TACACAGAG GAGCCAGAG TTTGTGTGTT AGGCTGAGT TTTGTGTGTT AGGCTGAGT TTTGTGTGTT AGGCTGAGT
6801 GAGGCTTGG TACACAGAG CCACTGCTCT TTTGTGTGTT AGGCTGAGT TTTGTGTGTT AGGCTGAGT TTTGTGTGTT AGGCTGAGT TTTGTGTGTT
CTCCGAGCC CTTGAGTCTT CTTGAGTCTT CTTGAGTCTT CTTGAGTCTT CTTGAGTCTT CTTGAGTCTT CTTGAGTCTT CTTGAGTCTT
6901 ATCAGAGT CAGAGCTCT CTTTGAAGC ACTAGAGAGC CCGAGAGT GGTGAGTCTT TTTGAGCAGA GAAGGTTAGA GTTAGGTTCTT CTTCTCTCTT
TACGCTCTCA GTCTCTCAAA AGTCTCTTTC TCAAGTCAAA AGTCTCTTTC TCAAGTCAAA AGTCTCTTTC TCAAGTCAAA AGTCTCTTTC
7001 CTAAGAGAGA GACTCTCTCT TTAAGAGAG AATGCTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC
GCTCTCTCTT CTAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC
7101 TTAATGCTAA AATGCTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC
GCTCTCTCTT CTAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC TCAAGTCTCC

FIG. 4B-2

10801 GTCTGAAA AAAAAGAGA GAGAGGAGG TGAGAGCCA AUAATCTDA CATTCTGTG GTTGTCTTG CTGTAGTCA TTCTGTAAG CAATGCTGC
CAGAGCTTT TTTTCTCT CTCTGCTTC ACTCTGGGT TATTAGATT GTTAGACAC CAACAGAAC GACATCAGAT AAGACTATTC GTTAGAGCG
10901 TTGCTGCCA GGTAGGAAGT AACATTCTT TATTAAGGT ATTGCTCTG CTTTATTTT CTGTTTATT TATGCTCTG AGATGAGAC CAGGAGCCT
AAGAGGGTT CCACTCTCA TTGTAAAGA ATATTTTCA TAAAGAGAC GAATAAAAA GACAAAATA ATACACGAC TCTTACTTG GTTCTGGA
11001 TGGCAAGCA GGTAGCTGT TTAGACTGA GCAATCTCC AGCTTGCAC TGGGGATTC TAGGCAAGG TTCTACCACT GAGCCACACT CCCCACCCC
AAGTTGGTT CCAATGACA AATGTTGACT CGGTATGAG TCGAAGCTG ACGGCTTAG ATCGTTTCC AAGATGATGA CTGGTGTGA GGGGTGGGG
11101 ATCCCTCTT GGAAGATTCT AGCCAGTTC ATACCTAGC TTGATCTTT TAGACGGTC TTAGTAGAC TCAATT
TAGGAGAGA CTTCTAGA TCGTCAGG TATGATCGG AACTAGAA ATCTGCCAG ATGATCTCG AGTCA

FIG. 4B-4




FIG. 5A






FIG. 5B

VEGF project	VEGFR2 project	Tie2 project
Screening primers	Screening primers	Screening primers
Primers: VF1-VR1A Product size: 0.69Kb	Primers: KF1-KR1 Product size: 0.45Kb	Primers: TF3-TR1 Product size: 0.45Kb
PCR program	PCR program	PCR program
Hot start	Hot start	Hot start
94°C 40 sec 65°C 1 min 30 sec 72°C 1 min 30 sec	94°C 40 sec 58°C 1 min 30 sec 72°C 1 min 30 sec	94°C 40 sec 58°C 1 min 30 sec 72°C 1 min 30 sec
40 cycles	40 cycles	40 cycles
Confirmation primers	Confirmation primers	Confirmation primers
Primers: VF2-VR2 Product size: 0.98Kb	Primers: KF2-KR2 Product size: 0.58Kb	Primers: TF2-TR1 Product size: 0.47Kb
PCR program	PCR program	PCR program
Hot start	Hot start	Hot start
94°C 40 sec 65°C 1 min 30 sec 72°C 1 min 30 sec	94°C 40 sec 65°C 1 min 30 sec 72°C 1 min 30 sec	94°C 40 sec 58°C 1 min 30 sec 72°C 1 min 30 sec
40 cycles	40 cycles	40 cycles

FIG. 6

**FIG.7**

*FIG.8*





FIG.9

**FIG.10**

SEQUENCE LISTING

<110> XENOVEN CORPORATION

<120> TARGETING CONSTRUCTS AND TRANSGENIC ANIMALS PRODUCED
THEREWITH

<130> PXE-008.PC

<140>

<141> 1999-12-16

<150> 60/152,522

<151> 1999-09-03

<160> 39

<170> PatentIn Ver. 2.0

<210> 1

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer PGKF

<400> 1

atcgaattct accgggtagg ggaggcgctt t

31

<210> 2

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer PGKR

<400> 2

ggctgcaggt cgaaaggccc ggagatgagg

30

<210> 3

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer NeoF

<400> 3

acctgcagcc aatatgggat cggccattga ac

32

<210> 4

<211> 37

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer NeoR

<400> 4

ggatccgcgg ccgccccag ctggttcttt ccgcctc

37

<210> 5
<211> 30
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer TKF

<400> 5
ggatcctcta gagtcgagca gtgtggtttt 30

<210> 6
<211> 30
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer TKR

<400> 6
gagctcccgt agtcaggttt agttcgccg 30

<210> 7
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer F5R51

<400> 7
gtacatttaa atcctgcagg 20

<210> 8
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer F5R52

<400> 8
agtcctgca ggatttaa 20

<210> 9
<211> 77
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer F3R31

<400> 9
ggcccggtt taattaatgc atcatatggt accgtttaaa cgcggccgca agcttgtcga 60
cggcgccg gccggcc 77

<210> 10
<211> 77
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer F3R32

<400> 10

gatcgccgg ccggcgccg gtcgacaagc ttgcggccgc gtttaaaccg taccatatga 60
tgcattaatt aagcccg 77

<210> 11

<211> 39

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer VN1R

<400> 11

ctgtatttaa atctgccac cctattcagg acagtagtc 39

<210> 12

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer VN1F

<400> 12

ccaatgcac aaccagcca ggaggagtgc g 31

<210> 13

<211> 38

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer VN2R

<400> 13

aacgcgtcga cttcggagat gtttcgggga taaccagg 38

<210> 14

<211> 40

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer VN2F

<400> 14

ttggcgccg ccataagaa gagacaccaa aggcacgctc 40

<210> 15

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer FosB1F

<400> 15

ctgtatttaa atcccgtttc tcaactgtgcc tgtgtc 36

<210> 16
<211> 35
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer FosB1R

<400> 16
gtctcctgca ggcttcctcc tccttggtcc ttgcg 35

<210> 17
<211> 35
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer FosB2F

<400> 17
aacgcgtcga cggatgggat tgacccccag ccctc 35

<210> 18
<211> 33
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer FosB2R

<400> 18
ttggcgcgcc ccttgctcc acctctcaaa tgc 33

<210> 19
<211> 31
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer VF1

<400> 19
acctcactct cctgtctccc ctgattccca a 31

<210> 20
<211> 25
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer VR1A

<400> 20
gctctggcgg tcacccccaa aagca 25

<210> 21
<211> 28
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer VF2

<400> 21
ccctttccaa gacccgtgcc atttgagc 28

<210> 22
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer VR2

<400> 22
actttgcccc tgtccctctc tctgttcgc 29

<210> 23
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer KF1

<400> 23
gctgcgtcca gatttgctct cagatgcg 28

<210> 24
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer KR1

<400> 24
ttctcaggca cagactcctt ctccgtccct 30

<210> 25
<211> 32
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer KF2

<400> 25
cagatggacg agaaaacagt agaggcgttg gc 32

<210> 26
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer KR2

<400> 26
gaggactcag ggcagaaaga gagcg 25

<210> 27
<211> 29
<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer TF3

<400> 27

agcttagcct gcaaggggtgg tctcatcg

29

<210> 28

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer TF2

<400> 28

caaatgcacc ccagagaaca gcttagcctg c

31

<210> 29

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer TR1

<400> 29

gctttcaaca actcacaact ttgcgacttc ccg

33

<210> 30

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer F51

<400> 30

cccagtgtct ctgatttagg gagagcacct gag

33

<210> 31

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer R51

<400> 31

ccagactgcc ttgggaaaag cgcctc

26

<210> 32

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer F52

<400> 32

cagtgagagt cttctctgtc cctcaatcgg ttctg

35

<210> 33
<211> 26
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer R52

<400> 33
tggatgtgga atgtgtgcca ggccag

26

<210> 34
<211> 32
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer F31

<400> 34
aatcaaagag gccaactgtg tgtgagaggt cc

32

<210> 35
<211> 26
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer R31

<400> 35
cggctcccca aaatgtggaa gcaagc

26

<210> 36
<211> 30
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer F32

<400> 36
gaatccatct tgctccaaca cccaacatc

30

<210> 37
<211> 26
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer R32

<400> 37
cgctctctct cccagtcctc cccttg

26

<210> 38
<211> 4998
<212> DNA
<213> Mus sp.

<400> 38

tccaccacc	tgtttctcac	gtccccggcc	ttcctagtta	acttcatggt	taaagaagcc	60
tcaccgggg	aggggtggt	gccacagaag	gaagggtgct	cccacaagcc	cccagtgctc	120
ctgatttagg	gagagcacct	gagcccagtg	agagtcttct	ctgtccctca	atcggttctg	180
aaattcccca	cttgccctcc	ttatccaggg	gacagggtcg	cccaccctat	tcaggacagt	240
agtcttaaac	tcgtagccaa	cagacttttt	attgggctgg	gagaaagaga	tgaggctcct	300
gaagctcagc	cgagtgggct	ctgattccta	cttctcagag	gtcgggcagc	ccagccaata	360
ctgagcaatg	gagcgtgggt	agggaggatt	cacagagtcc	actcgccggg	ttctaagggt	420
gactcggtag	tatttgtctg	aaagaaagaa	tggaaaaagg	gttatgtgag	attctgcctg	480
atcctgtcca	ctgggtccca	gaaggataaa	ggctttttct	cagaagggaa	agtgaacatc	540
caccaagcag	ataatgtcac	catctacagg	ctgtgttcag	caccagggga	ccaagacctg	600
caggcaaggc	ctagccaaaa	ccagtctaag	gagtagaaag	gggctcccac	ctccagagaa	660
gaaatagacg	ctctgaatgg	gctcgcaggt	ggcaggtaca	agccagtcga	tatcataatc	720
atagttgttg	taggttccta	gcccactctc	ctcgtctggag	aacaaagaga	accagattga	780
acgtgatgaa	cgacgggagt	tcgagctctg	gctgcgtctg	tggccacgcc	ctcggcgtga	840
acgatagcgc	tttcggcttc	tacgcttaga	cttctgtttt	ttggcttggg	cagagtggga	900
taaggagcca	gtgacgtaga	tgcggccggc	catagcagcg	tccactttcc	ctggcacacc	960
atgccagttc	cggctgatga	attgggggtc	tctggctcca	tctgtaacag	ggaagggggt	1020
aatgcacttg	gcagattctg	gctttgattt	ctccagcaag	gttgtctgtc	tatctattta	1080
tctatcttta	tctatgtatc	tatctatata	tctatgtatc	tatctatcta	tcactacct	1140
acctacttac	ctatctatgt	atctatctat	ctatcatcta	cctacctact	tacctatcta	1200
cctattttatt	tgtttgtttg	ttttctttga	aacaggatct	tagcacctac	ctatggctgg	1260
tttgcaactc	actatgaagc	cataactggc	ctcttaactc	acaaagatcc	acttgccctg	1320
gtctctgagt	gctgggatta	aaagcatgtg	ccactacacc	cagctccagt	aggaccttta	1380
gaacacattt	gctatgcctt	gcctaagaca	cacaactcag	tccccaggcc	ccagcctccc	1440
tgtctagagc	tttttcccat	cctctctcca	ctgtatccct	tgaatctctg	ccccatccga	1500
aaccctcag	cgcgcagccc	ctccttctgc	tgtgttaggc	aaagtccaag	gtatgggatc	1560
caaatagagc	caagcctcat	ccccaaaag	tcaacagaag	caaagtctag	ccagagcaaa	1620
cagctcttga	tcgatgggtg	cacagttcca	ggccctccc	ctggaagccc	ccactatcac	1680
agcccagttt	ccagagaaaag	aagccagcct	tgtctccct	ccataccaga	ggatctgccc	1740
cagaagagga	gttcgaaaat	gttctcccag	ctgtcccgt	gaagcaaggc	aaagtgtca	1800
aacacggctg	acagagagct	gccttcgcac	tcctcctggc	tgggttgctg	ctgaaattcg	1860
tactcccagt	actgcttccc	tgaggagcag	aacagctggc	atcaggagag	atctgaccaa	1920
ggcagagagg	aatcatggaa	tagaacaggg	actccaccac	ctgccccctt	ctcctccacc	1980
ctgagtaccc	ttgaagaagt	agaccctttc	cgggccactg	taacgggtggg	caggaagggc	2040
gaacgctgca	tcaacattgt	ctggtatgcc	actgaagcct	tcggagatgt	ttcggggata	2100
accaggggtc	aggaccccat	cctcaaagcg	ccagtactga	ctaccctgaa	agacagagat	2160
cagaaggggtg	aggacatacc	gctggccaca	gaagcagtc	tatatcctaa	actggctgtc	2220
acctgctcct	ggagtccctg	actgctttgt	cttcacagct	ccccagcacg	tccatggcac	2280
cctttacctt	gcctcagact	taggtctggt	accttgaaca	agtaggtctt	cccctgacag	2340
ttgatgcgag	tgaaggcagc	atcgatgggg	ccctcaatgc	cccagacatc	ttggataagt	2400
ttggggatcc	caggcctcac	tgccgtctca	tctagctcat	agcagtactg	ccctagaaca	2460
ggggaaactg	tgtgagaagc	agatgagcct	aaggcagatc	cgaccgccac	cagacctgtc	2520
catagagtca	cctcgggaag	caaagagggg	cccattcttg	agatccgtga	aggcgtcaaa	2580
gggctttcca	ctgcacagtt	cttcctctgg	aaactcaggg	gtcccttgat	cagtgggtgtc	2640
gggccttagg	atctcctcct	gttgctccac	tttaggcgct	gggggtgctg	gctgttcctc	2700
aggatctagg	aaggctgtcg	gcttttagagt	gccgtccgtc	cgaggattta	ggtcaccggg	2760
tggagaggtg	ttctcgggtt	gcacaccggt	gttggtattg	ttcttgggct	cctccacgta	2820
gtcatagctc	caataatcat	cctctggcat	agtgaacacg	tccccccgcg	ttactgcagg	2880
cagaacgggg	agcagtgagt	gtcaggctgt	ggaggggagc	ccaggcccac	ccaccagggc	2940
tctgaactca	ccttggggct	tgactgtctc	catgtagtgc	gcacagcagc	tctgatagta	3000
agtgcaaagc	tcgtcacact	gacacttctt	gctggccatg	aaacctgag	tgcagcggcc	3060
ccttgcatgac	tctatgggag	ggaatatcag	gctttacagc	caatctaggg	cacctgcccc	3120
acctgcactt	ccctaggtac	ccaccaatcc	cctcccacac	cttggtcagc	cagagaaacc	3180
catgccacca	gggctagtat	gaaaaagggc	ctcaggggtg	ccatggcgag	cctctagccc	3240
agggccttgg	caagctgggc	gcggagcttc	tggaaatctg	ctgtcctgcc	tgaaaaaaga	3300
agcagactga	agaagagttc	ctagttccct	gggtttctgc	cctttatttg	ctcatcctct	3360
ggcccagccc	cattgccttc	ctccaaacac	agctgcagca	aagggtcaca	ttcccagaac	3420
cccagcccca	ggagagctgg	gaaacagaaa	accctcgcca	agaccaaagt	cagtagggtc	3480
acgggcagga	gggataacac	gcttagctta	gctggggagg	tggaaagaag	catgtgttgt	3540
caccctctga	gccagtcctg	ttaatctccc	tgagccttac	tttttataaa	gtgggaccat	3600
ggtgccttgc	ctcatcaggt	gttgagagat	tccgtgagct	agaacagaca	aaacgtttcg	3660
tgcttgaggt	agcttccaac	tcatcccat	aagccgttat	cgatttactg	tttgatcagg	3720

```

ctagggtgctt gtcccatcct accccccgct tccaatctgg atttttgggg caagaagggg 3780
gggtggggga gagctggcaa gcactttggg ggaggttttc tttcttctc ataaaagaac 3840
aaagcttcat ttctggcctc tccttgttct ctctaagctg ggtgttacag cataggaagt 3900
agtgggtcag agtctattct tctttcttta ttttttttag atttatttat tttatgtttt 3960
gtgtataagt gtctgtcac atgtgcact gtgcaccaca tgcattgtct gtgtctatgg 4020
aggtcagaag agggctttga ataccctgga actggagttt tgaacagtta tgagctgccg 4080
tgtggatgct gagaatcaaa cccaggctct ctgtaagaac aagtactctt aaaggctgag 4140
ccatctttcc agtcccagag cccattcctg aggttttcac taatccattg atcctcgggg 4200
gaccaccctg gccacacctt caatgacctc atttatttta aaaaaaaat' ggactcattg 4260
ggcatacttt ctagactcac atactaagt ggattttctc ataaagaagt gctcactggg 4320
gtagagtgc aggttttggg ccaattcca agcactggca cacttctgaa gcccctccgt 4380
tttctgttct gtaatcacag gcgagcgtgc ctttgggtgc tcttctctat ggaccgcagt 4440
agtctcagcg gcaaaatgaa acactaaatt ttactcccta cagacgcgtg aagcctaagt 4500
ggaaaccggc attaaagggc tttaagaatc tcaactgcca ttctttaacc atccggaggg 4560
gacgtggata catgtagcca gcttgcttcc acattttggg gagccgagcg agcggtagga 4620
aatggaagac agctctttac agccctttct acagcatctt gcacaccacc aaggggagac 4680
tggggagagg aggcggagcc aggtgtgggc gtggctggag acctggggta ggcttgcgc 4740
tgctcgggg gcggagcccg tgaaacctag aggcggggcg tcaaatcctt gactctgctg 4800
ctcagaggcg tggttgctgt tgagcatctt agctccgctg tgcttagatt ggagcagcgc 4860
tttgttccgg gcaccggcgt ctctaccctc ccgctctgg tccatgcttc tctctccctt 4920
catgcccttc ctaagtgcgt gagtcccgga gctgccctcc tcttctgct tctacacttg 4980
tagcccgaca cctttacc
4998

```

<210> 39

<211> 11176

<212> DNA

<213> Mus sp.

<400> 39

```

gcagctgggc aaacgttggc gatgccggtg caaagtatat acccgggtggt tagcagaagc 60
tgagaacttt tagccgaaag ccggctccct aagccgaagc taggcaagta ggggaagaaa 120
aagaaaaaaa aaattccaga gaagcttcca gagcctcctc ctcttccctc ttcttcaaa 180
aggactgcaa gtccgcagtc accctccacc cagcaagagt tagggcctcg aacccgggtc 240
acgctgcctc cgcctcctgc cgaacgtaac gggggaccgc tgcgtaaaagc gtgacgcgct 300
ggaaatcctc gctctgacgc gggcacgcac aggcgcgagc cctccgccc gccccgccc 360
tgacgtccgg gcacgttcta ttttggaaag ccgaggccac gttgctaagg gagggggcag 420
cgtggctttg tgattggctg tcgcgcgcag ctttagccaa tcagcgttcc ctctctattt 480
gtagagcgta gctcccttcc ttgcttttgg tggttcttcc cgtgctgggg gtctccaaga 540
ggagagctag gattcttgtc gcgatcggga ctcgttgtca ccccatggtc tgcgaggact 600
tgtgtggacc tggctctgtg tcataagcta gaggttttgg gctgagtgtt agcgcctcta 660
agggggaact gaaggcctca tcttctcag gcacacatat acgtgctcct gagctctaga 720
cactcagtc ttccgaggtg ttcaaact agatgagcta gcctacggag aggcagccag 780
gtggtctcta aaaggtctgc ctcccttag ttccaggct ctgattggcc agggattcag 840
cccttccctc gccacgcccc ctagagtagt taagcctcta ggattccact tgcgggaagg 900
gggggggggg gggcgtgatg gacgttctt ggggacgcag atcctatgtc accccatccc 960
ctgcaagaca gtctgagaga ttctcgctgt cacttttctc tgcctatcag ttactgaaa 1020
cctgtcagtc tactgggaa gagacagaca ctcggaagg atgctctcaa ctcttaggcc 1080
ggtcccccaa caccgttggg actgggatct ccgctgagg gagccctcat gcagtggggg 1140
gtgtgtttgt gtgtgagtgg agaggaaggc ttggctaagg cctctccctc tccctccctc 1200
tgtgttgggg gttggggggg tttggctgta tgtgtgtgtg aatgtctgtg gctccatccc 1260
gggagtttgt caccaggttc tgtccagcct cctctcccac ccaccccc acacctaga 1320
gtcaccaacc cggggtgtga ttcaacccc gctggaaccg tgcaaccttt ccccgaggaa 1380
gaaggaggag gtagaaggca gttgaacaga atctctcatt aaccactgcg tcacggtgta 1440
gtggaagggt ggggtgtgtg gctttttgcc tgtgacacac acatccacac ccgctcacc 1500
tgtgtcact cacagggtcg gtctctctta tctctcttgg gcgtgtgtgt gtcgggtggc 1560
ttgttttgt gtctacgct gtgtgtgtat gtctcaccgc gtaggagtgc gccggtctcg 1620
gggaaatgcc cggctccttc gtgccaacgg tcaccgcaat cacaaccagc caggatcttc 1680
agtggctcgt gcaaccacc ctcatctctt ccattggcca gtcccagggg cagccactgg 1740
cctcccagcc tccagctgtt gacccttatg acatgccagg aaccagctac tcaaccccag 1800
gcctgagtgc ctacagcact ggcggggcaa gcggaagtgg tgggccttca accagcaca 1860
ccaccagtgg acctgtgtct gccggtccag ccagagccag gcctagaaga ccccgagaag 1920
agacagtaag tatgaggcct caggagttgg gatggaggag cctagctagg gatgtgggct 1980

```

cagtttgtac	agtgccttgc	tgccatgcat	gaagatccct	agcacagcat	aagccaggag	2040
tggttatgca	gacctgtaac	cccagctctc	agaaggtgga	ggcaggagga	gcaggagtgc	2100
gaggccagcc	tgtgctactt	atggagtcca	gcctgcactg	caagagatca	ttattttcaa	2160
aagttggcct	tggtgggagg	tggttgaggg	aagtaagaga	aagtgcagct	aatttttgtca	2220
cttaatatgtt	ggaggttcct	ctgaggcctc	aagtctgaag	gaactttacc	attctggcca	2280
gtgaggagta	gggtttatta	tttggggttc	aggaggaagg	aagttttctt	agggctgata	2340
gaggtacccc	cagatctcat	ggtccttata	tctgactcag	cttaccaccag	aagaagaaga	2400
aaagcgaagg	gttcgcagag	agcggaaaca	gctggctgca	gctaagtgca	ggaaccgtcg	2460
gagggagctg	acagatcgac	ttcaggcggt	aaggaggagt	ctgggggtgt	cttgaggccg	2520
tgctggggagc	actctgcctt	gttcttcccc	cgtttctcac	tgtgcctgtg	tcctaaacga	2580
ggaaaccccc	tcttagggaa	caggggtcag	tataggctga	tggagtggct	ccatatgcat	2640
gctcagaccc	atgccactt	actttcgact	gttccccact	ttccctgaat	atgtccccac	2700
atgtcacccct	cctggctttc	tctcagccta	aggagacaag	ctagaggagg	taattctctc	2760
acottctttt	cttactaaa	taataatcca	ttttgccttc	ctgcctccat	ttttttttcc	2820
tgagctgggg	atctacctgt	cgtagttcag	ccctcctccc	ccaacttgat	agcctcaagt	2880
ttcagccctt	ggctgagatg	ccatcatcct	gactggctct	ggctggaaac	tattttgtgc	2940
taagtcaatt	ccttgtctgc	tacttcagct	atctacagty	ctgccgaact	tgagctggtg	3000
gcgcccacca	agcccacttc	tttctctctt	ttttacctca	gtgcaacccc	ccacacacaa	3060
aacttcatgc	ctgccccctg	aaaccagggt	cgctctctga	ctccccgtcg	ggaggctgaa	3120
ggagatgggt	aacagaacct	cattaaaaac	aacacataag	cattacctac	tgactcaaca	3180
aactgtagtg	tttttctttt	ttcctctcaa	aaaattattt	cgtttgttta	tttattattt	3240
gottatgttt	gagtgagtgc	tggtgcacca	cagcacacat	acgaggtcag	agggaaattt	3300
tcatagtttg	ttctctcctt	ccgtgttgtg	ggtgcttgct	ggcaatctcc	ttcactcagt	3360
gagctacaat	gcccccttct	gcccccttaag	gcagagtact	ccttagtaca	gggggaccct	3420
ttcctcgccc	tctcaaagtt	gagattacaa	atgttcacca	tcacaccagg	cttgaggttc	3480
ttgcctatca	gtgacgtcca	ctcctgccta	gcttcttccc	aaccatcttt	tagtctgatg	3540
gggaaaccga	ggcacgagta	gcatggtcta	ccaggatttc	ctcttagggg	acggtccctt	3600
cagttggggag	ggagctgtcc	agccccctgg	atcagcagca	agaatgtatg	agtgtggggg	3660
tggggcgggtg	aagctactct	gtgtgggtcg	tgaccagcaa	ttctcctttc	tctgtctcct	3720
atgacctggc	cctgctggga	tccattagga	aactgatcag	cttgaagagg	aaaaggcaga	3780
gctggagtgc	gagatcgccg	agctgcaaaa	agagaaggaa	cgcctggagt	ttgtcctggg	3840
ggcccacaaa	ccgggctgca	agatccccta	cgaagagggg	ccggggccag	gcccgtggc	3900
cgaggtgaga	gatttgccag	ggtcaacatc	cgctaaggaa	gacggcttcg	gctggctgct	3960
gccgccccct	ccaccaccgc	ccctgccctt	ccagagcagc	cgagacgcac	cccccaacct	4020
gacggcttct	ctctttacac	acagtgaagt	tcaagtcttc	ggcgaccctt	tccccgttgt	4080
tagccttctg	tacacttctt	cgtttgctct	cacctgcccc	gaggtctccg	cgttcgccgg	4140
cgcccaacgc	accagcggca	gcgagcagcc	gtccgacccg	ctgaactcgc	cctcccttct	4200
tgctctgtaa	actctttaga	caaacaaaac	aaacaaaacc	gcaaggaaaca	aggaggagga	4260
agatgaggag	gagaggggag	gaagcagtc	gggggtgtgt	gtgtggaccc	tttgactctt	4320
ctgtctgacc	acctgccgcc	tctgccatcg	gacatgacgg	aaggacctcc	tttgtgtttt	4380
gtgtctgttc	tctggttttc	tgtgccccgg	cgagaccgga	gagctggtga	ctttggggac	4440
aggggggtggg	gcggggatga	acaccctctc	tgcatactct	tgtcctgtta	cttcaacca	4500
acttctgggg	atagatggct	gactgggtgg	gtagggtggg	gtgcaacgcc	cacctttggc	4560
gtcttacgtg	aggctggagg	ggaaagagtg	ctgagtgtgg	ggtgcagggt	gggttgaggt	4620
cgagctggca	tgcacctcca	gagagaccca	acgaggaaat	gacagcaccg	tcctgtcctt	4680
cttttcccc	accaccccat	ccaccctcaa	gggtgcaggg	tgaccaagat	agctctgttt	4740
tgctccctcg	ggccttagct	gattaactta	acatttccaa	gaggttacaa	cctcctcctg	4800
gacgaattga	gcccccgact	gagggaaagt	gatgccccct	ttgggagtct	gctaacccca	4860
cttcccgtcg	attccaaaat	gtgaaccctt	atctgactgc	tcagtctttc	cctcctggga	4920
aaactggctc	aggttggatt	tttttctctg	tctgctacag	agccccctcc	caactcaggc	4980
ccgtccccc	ccctgtgcag	tattatgcta	tgtccctctc	accctcacc	ccaccaccag	5040
cgcccttggc	cgtcctcggt	gggccttact	ggttttgggc	agcagggggc	gctgcgacgc	5100
ccactctgtc	ggagcgcttt	atactgtgaa	tgagtggctg	gattgtggg	cgcgccggat	5160
gggattgacc	cccagccctc	caaaaactttt	ctggggcctc	cccttcttcc	acttgcttcc	5220
tcctccctcc	tgacagggag	ttagactcga	aaggatgacc	acgacgcac	ccggtggcct	5280
tcttgctcag	gccccagact	ttttctcttt	aagtccctcg	ccttccccag	cctaggacgc	5340
caactctctc	ccaccctggg	agccccgcac	cctctcacag	aggtcgaggc	aattttcaga	5400
gaagttttca	gggctgaggc	tttggctccc	ctatcctcga	tatttgaatc	cccaaatagt	5460
ttttggacta	gcatacttaa	gagggggctg	agttccact	atccactctc	atccaattcc	5520
ttcagtccca	aagacgagtt	ctgtcccttc	cctccagctt	tcacctcgtg	agaatccac	5580
gagtcagatt	tctattttct	aatattgggg	agatgggccc	taccgcccgt	cccccgctg	5640
gcatggaaca	ttccataccc	tgctcctgggc	cctaggttcc	aaacctaata	ccaaacccca	5700

```

ccccagcta tttatccctt tccctgggtcc caaaaagcac ttatatctat tatgtataaa 5760
taaataatatt atatatgagt gtgcgtgtgt gtgcgtgtgc gtgcgtgcgt gcgtgcgtgc 5820
gagcttcctt gttttcaagt gtgctgtgga gttcaaaatc gcttctgggg atttgagtca 5880
gactttctgg ctgtcccttt ttgtcacttt ttgtgtgttg tctcggctcc tetggctgtt 5940
ggagacagtc ccggcctctc cctttatcct ttctcaagtc tgtctcgctc agaccacttc 6000
caacatgtct ccactctcaa tgactctgat ctccggtctg tctgttaatt ctggatttgt 6060
cggggacatg caattttact tctgtaagta agtgtgactg ggtggtagat tttttacaat 6120
ctatatcggt gagaattctg ggtggaaatg tctgatcagg agaagggcct gccactgccg 6180
accacaattc attgactcca tagccctcac ccaggctgta tttgtgattt ttttcatttt 6240
gtttttttgt attttgcacc tgaccccggt ggtgctgggg cagtctatca ctgggcagct 6300
ccctccccc ccttgggtct gcactgtgc caataaaaag cttttaaaaa actgtatcct 6360
tcagggtcaaa gtgtctgttt tccctggaca tctactacat ggcttccttt cagaaaaacg 6420
gagtttggat tgctagggaa gtcttgcctg cacttagtgg gacgcctaac gaatcagaac 6480
ctacaacggg actaaaagga agtggagact tgctaggttt tcccatgttc ccaggctggg 6540
ccacctactt gaaaaaataa ggggcggaag agtgaaggt accaaatttg gtgaagggtc 6600
tgggagaatt tcatgatcgg aaaagaattt attcaccttg ggtgtgcaat gaactttcag 6660
caacagttaa gggcaagggt gtaaaagctg ggcacaactt gtaaatccta gcatttgaga 6720
ggtggaggca aggggatcaa ctggtggagt tcagtgtcat gtggatcgta gataccaagc 6780
gcaaagatct gctatgggga gagggcttgg tacaccaggg gagccagaag tttcgtggtg 6840
agggtagtgg agggcaagtg gagagtgaga gttagcctca gggagattct acaggcaatg 6900
atgcagagtt cagacgctcc ctttgaaagc actagagagc cgcagcaggt tttgagcaga 6960
gaaggtaga gttaggtggt ctcttctagc ccatccagg ctgaggagga cgctgagggt 7020
ttcaagaagg atcgagaatg gaaagcagag gagaagaagg atccaagagg catggaggag 7080
gcagaacaca tttctcttct ttaatagcaa gcctggaaag gataacttgc tgcaggagga 7140
gatgctcacc agtcgggttg tctaggggtt tcttggaaaa gagaaggcat ttgctcaagc 7200
ctcggttccc ccattctcgc tcttctgtca gcttgtcttc cattaagtgt gtgtctcaag 7260
gccaccctgc tcaggactcc ttgtgagacg accttctatg ctcgagttca ttaaaaacac 7320
aattgcctgg tgccgtgctc tctccactgg ctcagttacc tcaaaagacc agggctaaag 7380
gtgtgatcac aactctatcc ccattactgc tccaacgcag agacaggact gagccggagt 7440
gaacaaatga acaaaaatga ctaataatgc atgctgtgatt aaatacataa aagagcagat 7500
gactggatga gcaaatcggt taaggagaga cagcaagatc ctagaatttt ggagactaat 7560
ttaaatccat ctttgagatg catttggtcg gaaattcctg ggaggaaaaa aagtgtaaat 7620
atgaagagag aataaatgag aataggggtg gcttcagaga ggtaactgc gcgtggtcg 7680
cttttgtaca agaatgtgaa ttgcaggag caaaatggga tagatactcc cgcccgaaag 7740
gtggaattga accactctgt cgctaaacag ctacaggttt gaagcctgca cccagacca 7800
ctaggatca tccgggcgaa aggagctatt tttagttagt tatataaagg cgagatacta 7860
ctacttttta cacttatggt cattatttgt ggtatcacgt agataattaa tttcaatggt 7920
ttcgaacatt ttttttcact ttttcttgtg aacatgtggt tcctcagtaa agtgttccgt 7980
gaatgactct actaactaaa aagtaagtag cttcatttgc atagcgcctt gcattttggg 8040
aagcagcgcc taaagtgcct gtctccctaa ctaaaagcag aattttttgc aaagtgtaaa 8100
gtcagtttta tttttgtttg tttgtttgct tgtttgttt taatggaaaa acttctcacg 8160
cggccattc gtagcagaat tctgagatttt ctgcaagcga gaagcaagac tttcgtaggg 8220
tctgacggca cgcggccgca gagcgacacc tgccgttgc ttatagaact gcaagtatgt 8280
agggaaatcta ctgagtcctt aggtgatgga gttgacaacc aactccctt gaggtttagac 8340
gctaaaaacc atcccttttt atatttatgt gattagccca gggaaactaa ggctcagaca 8400
tggaataatc cacagccgag ttcttgtagc ccaactccct aggggaaatg aaacctacag 8460
ttgtggtttt aatatgcttg gccaggggc agtggcccta ttggcaggag tggccttatt 8520
agcggagggt taccttggtt gagaagtgtg tcacttgagg gcgaggtttt gaggtacgta 8580
tgctcaagtc tggccagtgt gatcctggct gtctgcagaa cgtggtctcc ttctggtgc 8640
cttcggatca aggtgtagaa ctctcagctc cttctccagc accatgtctg cctgcttaat 8700
gctttgcttc tttccatgac gataatgaac tgtgcctctg aaactgtaag tcagccccc 8760
agttacatgt tttcttttat aagagttgca tatatatatg tatgtatata tgtatgtata 8820
tatgtatgta tatatatata tatatataaa cagggtctca ctcttagct ctggctggcc 8880
tgaaattcac tatatagccc aggattgcct gaactcttga gcaatcttcc tgcctcagc 8940
tcccaatggt attacaggca tgagtcacaa caagccattt aaatcttatg atgacttata 9000
agaagacaga aaatcagagt tcccttacct agttcacaga tccctacaat ctaacctcgt 9060
tcgctccata aacagcccta cccaccctc ctggaactgc tttgaggaaat gctgcaggct 9120
ctcacaggca cactcctcct tggttaatct cttcagcctg gttgccttcc ccccatgt 9180
ccatgtggcc caaagcctct catcctgttc tcaaatacca ctagctagta aggcctcccc 9240
acctgaccg gtttaaatat tagaaaaggg tcaactttct cctgccacag acaaccaaac 9300
caccatatgc ttgtcactta ctacctgact atgaagggtta atagatgtct tcacaacctt 9360
tctctgagcc tcagtttccc cacctgcata atgcatctga gacacagaat tccctagagc 9420

```

tgtggttctc	ctcattccta	gtgctgggac	cctttaatac	atttcctcat	gttgtggtga	9480
ccccaccacc	accataaaat	tatttccatt	gatacttcat	aactgtaatt	ttttctattg	9540
ttatgaatag	taatgtaagc	atgtgtgttt	cccagtgatc	ttagatgacc	ctgtggaaga	9600
gtcattccac	cccaaagggg	tccccaccac	aagttaagaa	ttcctgccat	agaggaatca	9660
cagggaccat	ggattaacac	ttgggtcgac	ttttgggctg	ccttctggga	ggcgctagag	9720
ctaatacacag	ctacatcaat	ttctgaaatt	ttgtgtgtgt	gtgtgtgtgt	gtgtgtgtgt	9780
gtgtgtgtgt	gccctgagtc	gggtgctgag	ataggccagt	ggcttttagtg	ttcctggacc	9840
cattactcac	cagaactctc	ccctcacctg	attctttgat	gtgaacacta	tgtcttcata	9900
gtggcggtgg	caatagcagc	aacagtgaac	taaattttta	aagtagaact	cagctggaga	9960
tacaaatatt	gcagttttga	agttgggggtg	gattgtctaa	taacttaata	acataaccca	10020
gaagagaggg	cccttgggtct	tgcaaacttt	atatgcctca	gtacagggga	acgccagggc	10080
caagaagtgg	gagtgggtgg	gtagggggagc	aggggtggggg	gagggtatag	gggactttcc	10140
ggatagcatt	tgaaatgtaa	atgaagaaaa	tatctaataa	aaatttgaaa	aaaaatgtta	10200
ccccagtttg	gcctggatct	cactacctca	accagactgg	catgtgactc	tgctgagatc	10260
tgccactctc	tgccctcctgg	gtgcagaaga	caatttttgg	aagttagttc	tcttcttcca	10320
tcttgtggat	tccagggatt	gaactcgggt	catcaggctt	ggctgcaagt	gacttactta	10380
ggtgtctccc	agaccctctc	ggtttgatta	gttagatgct	gcacttcattg	cctgactttc	10440
gcactatgta	gatagagcaa	tgtctataac	atctcctaca	atgatatgta	tatcaagagc	10500
caagtgatga	gatggctcag	tgggtaagag	cacagactgc	tcttccaaag	gtcccaggtt	10560
caaatcccag	caatcacata	gtggcttcca	ttccctctta	tggaatgtct	gaagactgct	10620
acagtgtact	tacatataat	aaataaataa	atcttaaaaa	aaaaaaaccc	agccgggctg	10680
ggtggcgcac	gcctttaatc	ccagcacttg	ggaggcagag	gcaggcggtt	tcctgagttc	10740
gacgccagcc	tggtctacag	agtgagttcc	acgacagcca	gaactacaca	gagaaaccct	10800
gtctcgaaaa	aaaaaagaga	gagaggggaag	tgagagcgca	ataatcttaa	catttctgtg	10860
gttgtctttg	ctgtagtcta	ttctgataag	caatgctggc	ttgctcccaa	ggtaggaagt	10920
aacatttctt	tataaaaggt	atgtgctctg	ctttattttt	ctgttttatt	tatggtgctg	10980
aggatggaac	ccaggaccct	tggcaagcaa	ggctagctgt	ttaccactga	gccatactcc	11040
agccttgcac	tgggggattc	taggcaaggg	ttctaccact	gagccacact	ccccacccc	11100
atccctctct	ggaagattct	aggcagttcc	atacctagcc	tttgatcttt	taagacggtc	11160
ttactagagc	tcagtt					11176

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 7 C12N15/90 A01K67/027

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
 IPC 7 C12N A01K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 30715 A (ISACOFF EHUD Y ;SIEGAL MICAH S (US); UNIV CALIFORNIA (US); CALIFOR) 16 July 1998 (1998-07-16) page 29, line 5 -page 30, line 10; claims ---	1-8,10, 22-26, 32,34,35
X	WO 98 28971 A (LINK CHRISTOPHER ;UNIV TECHNOLOGY CORP (US)) 9 July 1998 (1998-07-09) page 22, line 3 - line 6; figure 1 page 26, line 24 - line 28; claims 11-19; figure 1; example 3 page 18, line 9 - line 13 ---	1,2,8, 10,12, 14-17 1,2,7, 10-12, 14,17-20
Y	--- -/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

26 June 2000

Date of mailing of the international search report

03/07/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Chambonnet, F

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	CONTAG CH, ET AL.: "Visualizing gene expression in living mammals using a bioluminescent" PHOTOCHEMISTRY AND PHOTOBIOLOGY, (1997) VOL. 66, NO. 4, PP. 523-531. , XP000920570 the whole document	1,2,7, 10-12, 14,17-20
A	CONTAG PR, OLOMU IN, STEVENSON DK, CONTAG CH. : "Bioluminescent indicators in living mammals." NAT MED. , vol. 4, no. 2, February 1998 (1998-02), pages 245-247, XP000914588 the whole document	1
A	WO 98 36081 A (ANGELIS DINO A DE ;SLOAN KETTERING INST CANCER (US); MIESENBOCK GE) 20 August 1998 (1998-08-20) the whole document	1
A	WO 98 23633 A (CORNELL RES FOUNDATION INC) 4 June 1998 (1998-06-04) the whole document	1

Pat nt docum nt cited in search r port	Publication date	Patent family member(s)	Publication date
WO 9830715 A	16-07-1998	AU 5090498 A	03-08-1998
WO 9828971 A	09-07-1998	AU 5619298 A EP 0948612 A	31-07-1998 13-10-1999
WO 9836081 A	20-08-1998	EP 0981633 A	01-03-2000
WO 9823633 A	04-06-1998	AU 7410398 A	22-06-1998